

FORM PTO-1370  
(REV. 11-2000)

U.S. DEPARTMENT OF COMMERCE PATENT AND TRADEMARK OFFICE

ATTORNEY'S DOCKET NUMBER

TRANSMITTAL LETTER TO THE UNITED STATES  
DESIGNATED/ELECTED OFFICE (DO/EO/US)  
CONCERNING A FILING UNDER 35 U.S.C. 371

30434.4USWO

U.S. APPLICATION NO. (if known see 37 CFR 1.5)

09/869820

INTERNATIONAL APPLICATION NO.  
PCT/US00/00179INTERNATIONAL FILING DATE  
January 5, 2000PRIORITY DATE CLAIMED  
January 5, 1999

## TITLE OF INVENTION

METHODS FOR TREATING CONDITIONS ASSOCIATED WITH THE ACCUMULATION OF EXCESS EXTRACELLULAR MATRIX

## APPLICANT(S) FOR DO/EO/US

Nancy A. Noble, Wayne A. Border and Daniel A. Lawrence

Applicant herewith submits to the United States Designated/Elected Office (DO/EO/US) the following items and other information:

1. ☒ This is a **FIRST** submission of items concerning a filing under 35 U.S.C. 371.
2. ☐ This is a **SECOND** or **SUBSEQUENT** submission of items concerning a filing under 35 U.S.C. 371.
3. ☒ This is an express request to begin national examination procedures (35 U.S.C. 371(f)). The submission must include items (5), (6), (9) and (21) indicated below.
4. ☒ The US has been elected by the expiration of 19 months from the priority date (Article 31).
5. ☒ A copy of the International Application as filed (35 U.S.C. 371(c)(2))
  - a. ☒ is attached hereto (required only if not communicated by the International Bureau).
  - b. ☐ has been communicated by the International Bureau.
  - c. ☐ is not required, as the application was filed in the United States Receiving Office (RO/US).
6. ☐ An English language translation of the International Application as filed (35 U.S.C. 371(c)(2)).
  - a. ☐ is attached hereto.
  - b. ☐ has been previously submitted under 35 U.S.C. 154(d)(4).
7. ☒ Amendments to the claims of the International Application under PCT Article 19 (35 U.S.C. 371(c)(3))
  - a. ☒ are attached hereto (required only if not communicated by the International Bureau).
  - b. ☐ have been communicated by the International Bureau.
  - c. ☐ have not been made; however, the time limit for making such amendments has NOT expired.
  - d. ☐ have not been made and will not be made.
8. ☐ An English language translation of the amendments to the claims under PCT Article 19 (35 U.S.C. 371 (c)(3)).
9. ☒ An oath or declaration of the inventor(s) (35 U.S.C. 371(c)(4)).
10. ☐ An English language translation of the annexes of the International Preliminary Examination Report under PCT Article 36 (35 U.S.C. 371(c)(5)).

## Items 11 to 20 below concern document(s) or information included:

11. ☐ An Information Disclosure Statement under 37 CFR 1.97 and 1.98.
12. ☒ An assignment document for recording. A separate cover sheet in compliance with 37 CFR 3.28 and 3.31 is included.
13. ☐ A **FIRST** preliminary amendment.
14. ☐ A **SECOND** or **SUBSEQUENT** preliminary amendment.
15. ☐ A substitute specification.
16. ☐ A change of power of attorney and/or address letter.
17. ☐ A computer-readable form of the sequence listing in accordance with PCT Rule 13ter.2 and 35 U.S.C. 1.821 - 1.825.
18. ☒ A second copy of the published international application under 35 U.S.C. 154(d)(4).
19. ☐ A second copy of the English language translation of the international application under 35 U.S.C. 154(d)(4).
20. ☒ Other items or information: transmittal sheets, postcard, application data sheet

U.S. APPLICATION NO. (if known) <b>09/869820</b>		INTERNATIONAL APPLICATION NO. <b>PCT/US00/00179</b>		ATTORNEY'S DOCKET NUMBER <b>30434.4USWO</b>	
--	--	---	--	--	--

21. <input checked="" type="checkbox"/> The following fees are submitted: <b>BASIC NATIONAL FEE (37 CFR 1.492 (a) (1) - (5)):</b> Neither international preliminary examination fee (37 CFR 1.482) nor international search fee (37 CFR 1.445(a)(2)) paid to USPTO and International Search Report not prepared by the EPO or JPO. . . . . \$1000.00  International preliminary examination fee (37 CFR 1.482) not paid to USPTO but International Search Report prepared by the EPO or JPO . . . . . \$860.00  International preliminary examination fee (37 CFR 1.482) not paid to USPTO but international search fee (37 CFR 1.445(a)(2)) paid to USPTO . . . . . \$710.00  International preliminary examination fee (37 CFR 1.482) paid to USPTO but all claims did not satisfy provisions of PCT Article 33(1)-(4) . . . . . \$690.00  International preliminary examination fee (37 CFR 1.482) paid to USPTO and all claims satisfied provisions of PCT Article 33(1)-(4) . . . . . \$100.00  <b>ENTER APPROPRIATE BASIC FEE AMOUNT =</b>				<b>CALCULATIONS PTO USE ONLY</b>	
				\$	690.00
Surcharge of \$130.00 for furnishing the oath or declaration later than <input type="checkbox"/> 20 <input type="checkbox"/> 30 months from the earliest claimed priority date (37 CFR 1.492(e)).				\$	
CLAIMS	NUMBER FILED	NUMBER EXTRA	RATE	\$	
Total claims	55 - 20 =	35	x \$18.00	\$	630.00
Independent claims	9 - 3 =	6	x \$80.00	\$	480.00
MULTIPLE DEPENDENT CLAIM(S) (if applicable)				\$	0.00
<b>TOTAL OF ABOVE CALCULATIONS =</b>				\$	1,800.00
<input checked="" type="checkbox"/> Applicant claims small entity status. See 37 CFR 1.27. The fees indicated above are reduced by 1/2.				\$	900.00
<b>SUBTOTAL =</b>				\$	900.00
Processing fee of \$130.00 for furnishing the English translation later than <input type="checkbox"/> 20 <input type="checkbox"/> 30 months from the earliest claimed priority date (37 CFR 1.492(f)).				\$	0.00
<b>TOTAL NATIONAL FEE =</b>				\$	900.00
Fee for recording the enclosed assignment (37 CFR 1.21(h)). The assignment must be accompanied by an appropriate cover sheet (37 CFR 3.28, 3.31). \$40.00 per property +				\$	40.00
<b>TOTAL FEES ENCLOSED =</b>				\$	940.00
				Amount to be refunded:	\$
				charged:	\$

a. ☐ A check in the amount of \$ \_\_\_\_\_ to cover the above fees is enclosed.

b. ☒ Please charge my Deposit Account No. 50-0306 in the amount of \$ 940.00 to cover the above fees.  
 A duplicate copy of this sheet is enclosed.

c. ☒ The Commissioner is hereby authorized to charge any additional fees which may be required, or credit any  
 overpayment to Deposit Account No. 50-0306. A duplicate copy of this sheet is enclosed.


d. ☐ Fees are to be charged to a credit card. **WARNING:** Information on this form may become public. Credit card  
 information should not be included on this form. Provide credit card information and authorization on PTO-2038.

**NOTE:** Where an appropriate time limit under 37 CFR 1.494 or 1.495 has not been met, a petition to revive (37 CFR  
 1.137 (a) or (b)) must be filed and granted to restore the application to pending status.

SEND ALL CORRESPONDENCE TO

SaraLynn Mandel  
 Mandel & Adriano  
 35 N. Arroyo Parkway, Suite 60  
 Pasadena, California 91103

  
 SIGNATURE  
 SaraLynn Mandel  
 NAME  
 31,853  
 REGISTRATION NUMBER

In re application of UNIVERSITY OF UTAH et al.

International Application No.: PCT/US00/00179

International Filing Date: January 5, 2000

Title: METHODS FOR TREATING CONDITIONS ASSOCIATED WITH THE  
ACCUMULATION OF EXCESS EXTRACELLULAR MATRIX

Agent's Reference: 30434.4WO01

35 N. Arroyo Parkway

Pasadena, CA 91103

August 4, 2000

WIPO

34, chemin des Colombettes

CH-1211 Geneva 20

SWITZERLAND

and

Assistant Commissioner for Patents

BOX PCT

Washington, D.C. 20231

Dear Sir:

**COMMUNICATION**

Please consider the following amendments to the claims:

IN THE CLAIMS:

Please amend claim 32 as follows:

- 32. (Amended) A method for preventing or reducing excess extracellular matrix accumulation in a tissue or organ or at a wound site comprising inhibiting the overproduction of TGFB present in an organ or tissue or at a wound site to prevent the excess accumulation of extracellular matrix by administering a combination of agents that inhibit TGFB activity and/or production.--

In re application of UNIVERSITY OF UTAH et al.

International Application No.: PCT/US00/00179

International Filing Date: January 5, 2000

Title: METHODS FOR TREATING CONDITIONS ASSOCIATED WITH THE  
ACCUMULATION OF EXCESS EXTRACELLULAR MATRIX

Agent's Reference: 30434.4WO01

Please add new claims 52-55 as follows:

- 52. (New) A method for treating a condition associated with the excess accumulation of extracellular matrix in a tissue and/or organ or at a wound site by reducing the accumulation of extracellular matrix in the tissue and/or organ or at the wound site comprising administering at least one agent that degrades excess accumulated extracellular matrix.
- 53. (New) The method of claim 52, wherein said agent is a protease. —
- 54. (New) The method of claim 53, wherein said protease is plasmin.—
- 55. (New) The method of claim 52, further comprising the step of administering an agent or combination of agents that inhibit TGF $\beta$  activity and/or production.

#### REMARKS

The above amendments to the claims and the new claims are supported throughout the present specification, for example on page 10, lines 5-10, pages 19-20 and the original claims as filed. No new matter is added by the amendments or the new claims. Exhibit 1 includes the pending claims with the amendments indicated above.

Agent's Reference: 30434.4WO01

Sarah B. Allison

1941-1942  
1943-1944  
1945-1946  
1947-1948  
1949-1950  
1951-1952  
1953-1954  
1955-1956  
1957-1958  
1959-1960  
1961-1962  
1963-1964  
1965-1966  
1967-1968  
1969-1970  
1971-1972  
1973-1974  
1975-1976  
1977-1978  
1979-1980  
1981-1982  
1983-1984  
1985-1986  
1987-1988  
1989-1990  
1991-1992  
1993-1994  
1995-1996  
1997-1998  
1999-2000  
2001-2002  
2003-2004  
2005-2006  
2007-2008  
2009-2010  
2011-2012  
2013-2014  
2015-2016  
2017-2018  
2019-2020  
2021-2022  
2023-2024  
2025-2026  
2027-2028  
2029-2030  
2031-2032  
2033-2034  
2035-2036  
2037-2038  
2039-2040  
2041-2042  
2043-2044  
2045-2046  
2047-2048  
2049-2050  
2051-2052  
2053-2054  
2055-2056  
2057-2058  
2059-2060  
2061-2062  
2063-2064  
2065-2066  
2067-2068  
2069-2070  
2071-2072  
2073-2074  
2075-2076  
2077-2078  
2079-2080  
2081-2082  
2083-2084  
2085-2086  
2087-2088  
2089-2090  
2091-2092  
2093-2094  
2095-2096  
2097-2098  
2099-2100  
2101-2102  
2103-2104  
2105-2106  
2107-2108  
2109-2110  
2111-2112  
2113-2114  
2115-2116  
2117-2118  
2119-2120  
2121-2122  
2123-2124  
2125-2126  
2127-2128  
2129-2130  
2131-2132  
2133-2134  
2135-2136  
2137-2138  
2139-2140  
2141-2142  
2143-2144  
2145-2146  
2147-2148  
2149-2150  
2151-2152  
2153-2154  
2155-2156  
2157-2158  
2159-2160  
2161-2162  
2163-2164  
2165-2166  
2167-2168  
2169-2170  
2171-2172  
2173-2174  
2175-2176  
2177-2178  
2179-2180  
2181-2182  
2183-2184  
2185-2186  
2187-2188  
2189-2190  
2191-2192  
2193-2194  
2195-2196  
2197-2198  
2199-2200  
2201-2202  
2203-2204  
2205-2206  
2207-2208  
2209-2210  
2211-2212  
2213-2214  
2215-2216  
2217-2218  
2219-2220  
2221-2222  
2223-2224  
2225-2226  
2227-2228  
2229-2230  
2231-2232  
2233-2234  
2235-2236  
2237-2238  
2239-2240  
2241-2242  
2243-2244  
2245-2246  
2247-2248  
2249-2250  
2251-2252  
2253-2254  
2255-2256  
2257-2258  
2259-2260  
2261-2262  
2263-2264  
2265-2266  
2267-2268  
2269-2270  
2271-2272  
2273-2274  
2275-2276  
2277-2278  
2279-2280  
2281-2282  
2283-2284  
2285-2286  
2287-2288  
2289-2290  
2291-2292  
2293-2294  
2295-2296  
2297-2298  
2299-2300  
2301-2302  
2303-2304  
2305-2306  
2307-2308  
2309-2310  
2311-2312  
2313-2314  
2315-2316  
2317-2318  
2319-2320  
2321-2322  
2323-2324  
2325-2326  
2327-2328  
2329-2330  
2331-2332  
2333-2334  
2335-2336  
2337-2338  
2339-2340  
2341-2342  
2343-2344  
2345-2346  
2347-2348  
2349-2350  
2351-2352  
2353-2354  
2355-2356  
2357-2358  
2359-2360  
2361-2362  
2363-2364  
2365-2366  
2367-2368  
2369-2370  
2371-2372  
2373-2374  
2375-2376  
2377-2378  
2379-2380  
2381-2382  
2383-2384  
2385-2386  
2387-2388  
2389-2390  
2391-2392  
2393-2394  
2395-2396  
2397-2398  
2399-2400  
2401-2402  
2403-2404  
2405-2406  
2407-2408  
2409-2410  
2411-2412  
2413-2414  
2415-2416  
2417-2418  
2419-2420  
2421-2422  
2423-2424  
2425-2426  
2427-2428  
2429-2430  
2431-2432  
2433-2434  
2435-2436  
2437-2438  
2439-2440  
2441-2442  
2443-2444  
2445-2446  
2447-2448  
2449-2450  
2451-2452  
2453-2454  
2455-2456  
2457-2458  
2459-2460  
2461-2462  
2463-2464  
2465-2466  
2467-2468  
2469-2470  
2471-2472  
2473-2474  
2475-2476  
2477-2478  
2479-2480  
2481-2482  
2483-2484  
2485-2486  
2487-2488  
2489-2490  
2491-2492  
2493-2494  
2495-2496  
2497-2498  
2499-2500  
2501-2502  
2503-2504  
2505-2506  
2507-2508  
2509-2510  
2511-2512  
2513-2514  
2515-2516  
2517-2518  
2519-2520  
2521-2522  
2523-2524  
2525-2526  
2527-2528  
2529-2530  
2531-2532  
2533-2534  
2535-2536  
2537-2538  
2539-2540  
2541-2542  
2543-2544  
2545-2546  
2547-2548  
2549-2550  
2551-2552  
2553-2554  
2555-2556  
2557-2558  
2559-2560  
2561-2562  
2563-2564  
2565-2566  
2567-2568  
2569-2570  
2571-2572  
2573-2574  
2575-2576  
2577-2578  
2579-2580  
2581-2582  
2583-2584  
2585-2586  
2587-2588  
2589-2590  
2591-2592  
2593-2594  
2595-2596  
2597-2598  
2599-2600  
2601-2602  
2603-2604  
2605-2606  
2607-2608  
2609-2610  
2611-2612  
2613-2614  
2615-2616  
2617-2618  
2619-2620  
2621-2622  
2623-2624  
2625-2626  
2627-2628  
2629-2630  
2631-2632  
2633-2634  
2635-2636  
2637-2638  
2639-2640  
2641-2642  
2643-2644  
2645-2646  
2647-2648  
2649-2650  
2651-2652  
2653-2654  
2655-2656  
2657-2658  
2659-2660  
2661-2662  
2663-2664  
2665-2666  
2667-2668  
2669-2670  
2671-2672  
2673-2674  
2675-2676  
2677-2678  
2679-2680  
2681-2682  
2683-2684  
26

## Exhibit 1

1. A method for treating a condition associated with the excess accumulation of extracellular matrix in a tissue and/or organ or at a dermal wound site comprising:
  - a) reducing the excess accumulation of extracellular matrix associated with TGF $\beta$  overproduction and/or activity in an organ or tissue, or at a wound site; and
  - b) degrading excess accumulated extracellular matrix in said tissue and/or organ or wound site;whereby the accumulation of extracellular matrix in said tissue and/or organ or wound site is reduced from the level existing at the time of treatment.
2. The method of claim 1 wherein the accumulation of extracellular matrix is reduced to a level which does not interfere with normal functioning of the tissue or organ or result in scarring.
3. The method of claim 1 wherein said step of reducing the accumulation of extracellular matrix comprises administering at least one agent that inhibits TGF $\beta$  in an amount sufficient to inhibit TGF $\beta$  overproduction and/or activity.
4. The method of claim 3, wherein said agent that inhibits TGF $\beta$  is selected from the group consisting of selected from the group consisting of inhibitors of aldosterone, inhibitors of angiotensin II, anti-TGF $\beta$  antibodies, renin, ACE inhibitors, AII receptor antagonists, proteoglycans and ligands for the TGF $\beta$  receptor.
5. The method of claim 4 wherein said agent is a proteoglycan selected from the group consisting of decorin, biglycan, fibromodulin, lumican, betaglycan and endoglin.
6. The method of claim 4 wherein said ACE inhibitor is Enalapril<sup>tm</sup>.

7. The method of claim 4 wherein said AII receptor antagonist is Losartan<sup>tm</sup>.
8. The method of claim 1 wherein said step of reducing the accumulation of extracellular matrix associated with TGF $\beta$  activity comprises contacting renin with an anti-renin agent.
9. The method of claim 1 wherein said step of degrading excess accumulated extracellular matrix comprises contacting said matrix with at least one protease in an amount sufficient to degrade excess accumulated extracellular matrix to a level that does not impair the normal function of said tissue and/or organ or result in scarring.
10. The method of claim 9 wherein said protease is selected from the group consisting of serine proteases, metalloproteinases and protease combinations.
11. The method of claim 1 wherein said step of degrading accumulated extracellular matrix comprises administering an agent which increases the amount of active protease sufficient to degrade excess accumulated matrix to a level that does not impair the normal function of said tissue and/or organ or result in scarring.
12. The method of claim 11 wherein said protease is selected from the group consisting of serine proteases, metalloproteinases and protease combinations.
13. The method of claim 12 wherein said protease is plasmin and said agent which increases the amount of active plasmin is tPA.
14. The method of claim 12 wherein said protease is plasmin and said agent which increases the amount of active plasmin is a PAI-1 mutant.
15. The method of claim 1 wherein said condition associated with the excess accumulation of extracellular matrix is a fibrotic condition.

16. The method of claim 15 wherein said fibrotic condition is selected from the group consisting of glomerulonephritis, adult or acute respiratory distress syndrome (ARDS), diabetes, diabetic kidney disease, liver fibrosis, kidney fibrosis, lung fibrosis, post infarction cardiac fibrosis, fibrocystic diseases, fibrotic cancer, post myocardial infarction, left ventricular hypertrophy, pulmonary fibrosis, liver cirrhosis, veno-occlusive disease, post-spinal cord injury, post-retinal and glaucoma surgery, post-angioplasty restenosis, renal interstitial fibrosis, arteriovenous graft failure and scarring.
17. The method of claim 1 wherein said tissue or organ is selected from the group consisting of kidney, lung, liver, heart, arteries, skin and the central nervous system.
18. The method of claim 1 wherein said condition associated with the excess accumulation of extracellular matrix is scarring.
19. The method of claim 3 wherein said agent that inhibits TGF $\beta$  is nucleic acid encoding the agent.
20. The method of claim 9 wherein said protease is nucleic acid encoding a protease.
21. A method for treating a condition associated with the excess accumulation of extracellular matrix in a tissue and/or organ or at a wound site comprising:
  - a) administering at least one agent that inhibits TGF $\beta$  to reduce additional accumulation of extracellular matrix associated with TGF $\beta$  activity in a tissue and/or organ or at a wound site; and
  - b) administering at least one agent to degrade excess accumulated extracellular matrix; whereby the accumulation of extracellular matrix in said tissue and/or organ or at said wound site is reduced to a level which does not interfere with normal functioning of the tissue or organ in which said extracellular matrix accumulated.



22. The method of claim 21 wherein said agent that inhibits TGF $\beta$  and said agent to degrade excess accumulated extracellular matrix are administered concurrently.
23. The method of claim 21 wherein said agent that inhibits TGF $\beta$  and said agent to degrade excess accumulated extracellular matrix are administered sequentially.
24. The method of claim 21 wherein the agent that inhibits TGF $\beta$  is an anti-TGF $\beta$  specific agent that binds to and inhibits the activity of TGF $\beta$ .
25. The method of claim 21 wherein the agent that inhibits TGF $\beta$  is an agent selected from the group consisting of inhibitors of aldosterone, inhibitors of angiotensin II, anti-TGF $\beta$  antibodies, renin, ACE inhibitors, AII receptor antagonists, proteoglycans and ligands for the TGF $\beta$  receptor.
26. A method for treating or preventing a condition associated with the excess accumulation of extracellular matrix in a tissue and/or organ, or at a wound site, comprising administering a combination of agents in an amount sufficient to inhibit TGF $\beta$  activity and/or production to prevent or reduce the excess accumulation of extracellular matrix associated with TGF $\beta$  overproduction in a tissue and/or organ, or at a wound site.
27. The method of claim 26 wherein the agents to inhibit TGF $\beta$  comprise an agent that binds to and inhibits the activity of TGF $\beta$  and an agent selected from the group consisting of inhibitors of aldosterone, inhibitors of angiotensin II, inhibitors of renin, ACE inhibitors and AII receptor antagonists.
28. The method of claim 27 wherein said agent that binds to and inhibits the activity of TGF $\beta$  is selected from the group consisting of anti-TGF $\beta$  antibodies, proteoglycans and ligands for the TGF $\beta$  receptor.

29. The method of claim 28 wherein said agent that binds to and inhibits the activity of TGF $\beta$  is a proteoglycan selected from the group consisting of decorin, biglycan, fibromodulin, lumican, betaglycan and endoglin.
30. The method of claim 26 wherein said agents are nucleic acids encoding said respective agents.
31. The method of claim 26 wherein said agents are administered to reduce TGF $\beta$  overproduction prior to excess accumulation of extracellular matrix.
32. (Amended) A method for preventing or reducing excess extracellular matrix accumulation in a tissue or organ or at a wound site comprising inhibiting the overproduction of TGF $\beta$  present in an organ or tissue or at a wound site to prevent the excess accumulation of extracellular matrix by administering a combination of agents that inhibit TGF $\beta$  activity and/or production.
33. The method of claim 32 further comprising the step of degrading excess accumulated extracellular matrix in said tissue and/or organ or at said wound site.
34. The method of claim 32 wherein said accumulation of extracellular matrix is reduced to a level which does not interfere with the normal functioning of the tissue and/or organ in which said extracellular matrix accumulated or scarring is prevented or reduced.
35. A composition for reducing TGF $\beta$  overproduction in a tissue and/or organ or at a wound site comprising a combination of agents that inhibit TGF $\beta$  activity and/or production in a pharmaceutically acceptable carrier.
36. The composition of claim 35 wherein said agents comprise an agent that binds to and inhibits TGF $\beta$  and an agent that inhibits a pathway resulting in TGF $\beta$  production.

37. A composition for preventing or reducing excess accumulation of extracellular matrix associated with TGF $\beta$  activity comprising at least one agent that inhibits TGF $\beta$  activity and/or production and at least one agent that degrades excess accumulated extracellular matrix in a pharmaceutically acceptable carrier.
38. The composition of claim 37 wherein said agent that inhibits TGF $\beta$  is an agent that binds to and inhibits TGF $\beta$ , and said agent that degrades excess accumulated extracellular matrix is a protease.
39. The composition of claim 38 wherein said agent that inhibits TGF $\beta$  is selected from the group consisting of inhibitors of aldosterone, inhibitors of Angiotensin II, anti-TGF $\beta$  antibodies, renin, ACE inhibitors, AII receptor antagonists, proteoglycans and ligands for the TGF $\beta$  receptor.
40. The composition of claim 38 wherein said protease is selected from the group consisting of serine proteases, metalloproteinases and protease combinations.
41. The composition of claim 39 wherein said proteoglycan is selected from the group consisting of decorin, biglycan, fibromodulin, lumican, betaglycan and endoglin.
42. The composition of claim 37 wherein said agent that inhibits TGF $\beta$  and said agent that degrades excess accumulated extracellular matrix are nucleic acids encoding said agents respectively.
43. The composition of claim 37 wherein said agent that degrades excess accumulated extracellular matrix is an agent that increases the amount of active protease present in the tissue and/or organ or at a wound site.
44. The composition of claim 43 wherein said protease is plasmin and said agent is tPA.

45. The composition of claim 43 wherein said protease is plasmin and said agent is a PAI-1 mutant.
46. A composition for reducing the excess accumulation of extracellular matrix in a tissue and/or organ, or at a dermal wound site, comprising an agent that increases the amount of active protease present in the tissue and/or organ or at the wound site in a pharmaceutically acceptable carrier.
47. The composition of claim 46 wherein said protease is plasmin and said agent is tPA.
48. The composition of claim 46 wherein said protease is plasmin and said agent is a PAI-1 mutant.
49. A composition for treating a condition associated with the excess accumulation of extracellular matrix in a tissue and/or organ or at a dermal wound site comprising a combination of agents that inhibit TGF $\beta$  in a pharmaceutically acceptable carrier.
50. The composition of claim 49 wherein said agents that inhibit TGF $\beta$  are selected from the group consisting of inhibitors of aldosterone, inhibitors of inhibitors of aldosterone, inhibitors of angiotensin II, anti-TGF $\beta$  antibodies, renin, ACE inhibitors, AII receptor antagonists, proteoglycans and ligands for the TGF $\beta$  receptor.
51. The composition of claim 50 wherein said proteoglycans are selected from the group consisting of decorin, biglycan, fibromodulin, lumican, betaglycan and endoglin.
52. (New) A method for treating a condition associated with the excess accumulation of extracellular matrix in a tissue and/or organ or at a wound site by reducing the accumulation of extracellular matrix in the tissue and/or organ or at the wound site comprising administering at least one agent that degrades excess accumulated extracellular matrix.

53. (New) The method of claim 52, wherein said agent is a protease.
54. (New) The method of claim 53, wherein said protease is plasmin.
55. (New) The method of claim 52, further comprising the step of administering an agent or combination of agents that inhibit TGF $\beta$  activity and/or production.

**METHODS FOR TREATING CONDITIONS ASSOCIATED WITH THE  
ACCUMULATION OF EXCESS EXTRACELLULAR MATRIX**

By, Nancy A. Noble, Wayne A. Border and Daniel A. Lawrence

**FIELD OF THE INVENTION**

This invention relates to a method for preventing or reducing excess accumulation of extracellular matrix in tissues or organs or at a wound site, and more particularly to the prevention and treatment of conditions resulting from excess accumulation of extracellular matrix, using a combination of agents that inhibit TGF $\beta$ , or a combination of agents that inhibit TGF $\beta$  and agents that degrade excess accumulated extracellular matrix.

**BACKGROUND OF THE INVENTION**

Excess deposition and accumulation of extracellular matrix (ECM) is found in diseases such as fibrosis of the kidney or lung. Although the cytokine transforming growth factor Beta (TGF $\beta$ ) regulates extracellular matrix deposition for tissue repair, overproduction of TGF $\beta$  clearly underlies tissue fibrosis caused by excess deposition of extracellular matrix resulting in disease (Border and Ruoslahti, *J. Clin. Invest.* 90:1-7 (1992)). TGF $\beta$ 's fibrogenic action results from simultaneous stimulation of matrix protein synthesis (Border et al., *Kidney Int* 37:689-695 (1990), inhibition of matrix degradation and turnover and enhanced cell-matrix interactions through modulation of integrin receptors that facilitate ECM assembly. Overproduction of TGF $\beta$  has been demonstrated in glomerulonephritis (Okuda et al., *J. Clin. Invest.* 86:453-462 (1990)), diabetic nephropathy and hypertensive glomerular injury and in related fibrotic disorders of the lung, liver, heart, arterial wall, skin, brain, joints and bone marrow (Border and Noble, *N. Eng. J. Med.* 331:1286-1292 (1994)). In addition to the kidney, blocking the action of TGF $\beta$  with an agent such as antibody or the proteoglycan decorin has been shown to be therapeutic in fibrosis and scarring of the skin, lung, central nervous system and arterial wall (Border and Noble, *Kidney Int.* 51:1388-1396 (1997)).

Suppression of the production of ECM and prevention of excess accumulation of mesangial matrix in glomeruli of glomerulonephritic rats has been demonstrated by intravenous

administration of neutralizing antibodies specific for TGF $\beta$  (Border et al., *Nature* 346:371-374 (1990)) or administration of purified decorin, a proteoglycan (Border et al., *Nature* 360:361-364 (1992)) and by introduction of nucleic acid encoding decorin, a TGF $\beta$ -inhibitory agent, into a rat model of acute mesangial glomerulonephritis (Isaka et al., *Nature Med.* 2:418-423 (1996)).  
5 Inhibition of TGF $\beta$  activity, using for example anti-TGF $\beta$  antibodies, has been shown to disrupt TGF $\beta$  overproduction (Sharma et al., *Diabetes* 45:522-530 (1996)).

Dermal scarring following dermal injury results from excessive accumulation of fibrous tissue made up of collagen, fibronectin and proteoglycans at a wound site. Because the fibrous  
10 extracellular matrix lacks elasticity, scar tissue can impair essential tissue function as well as result in an undesirable cosmetic appearance. TGF $\beta$  is believed to induce the deposition of fibrous matrix at the wound site (Shah et al., *Lancet* 339:213-214 (1992)).

One explanation for persistent TGF $\beta$  overexpression in progressive fibrotic kidney disease is that repeated or multiple episodes of tissue injury, such as occurs in chronic diseases such as hypertension, diabetes or immune complex disease lead to continuous overproduction of TGF $\beta$  and extracellular matrix resulting in tissue fibrosis (See Border and Noble, *N. Eng. J. Med.* 331:1286-1292 (1994)). Another possible explanation for persistent TGF $\beta$  overexpression is the presence of a biologically complex interconnection between TGF $\beta$  and the renin-  
20 angiotensin system (RAS) in the kidney as part of an emergency system that responds to the threat of tissue injury as discussed further herein.

Renin is an aspartyl proteinase synthesized by juxtaglomerular kidney cells and mesangial cells in humans and rats. (Chansel et al., *Am. J. Physiol.* 252:F32-F38 (1987) and Dzau and  
25 Kreisberg, *J. Cardiovasc. Pharmacol.* 8(Suppl 10):S6-S10 (1986)). Renin plays a key role in the regulation of blood pressure and salt balance. Its major source in humans is the kidney where it is initially produced as prorenin. Signal peptide processing and glycosylation are followed by secretion of prorenin and its enzymatically active form, mature renin. The active enzyme triggers a proteolytic cascade by cleaving angiotensinogen to generate angiotensin I, which is in  
30 turn converted to the vasoactive hormone angiotensin II by angiotensin converting enzyme ("ACE").

The sequence of the human renin gene is known (GenBank entry M26901). Recombinant human renin has been synthesized and expressed in various expression systems (Sielecki et al., *Science* 243:1346-1351 (1988), Mathews et al., *Protein Expression and Purification* 7:81-91 (1996)). Inhibitors of renin's enzymatic site are known (Rahuel et al., *J. Struct. Biol.* 107:227-236 (1991); Badasso et al., *J. Mol. Biol.* 223:447-453 (1992); and Dhanaraj et al., *Nature* 357:466-472 (1992)) including an orally active renin inhibitor in primates, Ro 42-5892 (Fischli et al., *Hypertension* 18:22-31 (1991)). Renin-binding proteins and a cell surface renin receptor on human mesangial cells have been identified (Campbell and Valentijn, *J. Hypertens.* 12:879-890 (1994), Nguyen et al., *Kidney Internat.* 50:1897-1903 (1996) and Sealey et al., *Amer. J. Hyper.* 9:491-502 (1996)).

The renin-angiotensin system (RAS) is a prototypical systemic endocrine network whose actions in the kidney and adrenal glands regulate blood pressure, intravascular volume and electrolyte balance. In contrast, TGF $\beta$  is considered to be a prototypical cytokine, a peptide signaling molecule whose multiple actions on cells are mediated in a local or paracrine manner. Recent data however, indicate that there is an intact RAS in many tissues whose actions are entirely paracrine and TGF $\beta$  has wide-ranging systemic (endocrine) effects. Moreover, RAS and TGF $\beta$  act at various points to regulate the actions of one another.

In a systemic response to an injury such as a wound, the RAS rapidly generates AII that acts by vasoconstriction to maintain blood pressure and later stimulates the secretion of aldosterone, resulting in an increase in intravascular volume. In the wound, TGF $\beta$  is rapidly released by degranulating platelets and causes a number of effects including: 1) autoinduction of the production of TGF $\beta$  by local cells to amplify biological effects; 2) chemoattraction of monocyte/macrophages that debride and sterilize the wound and fibroblasts that begin synthesis of ECM; 3) causing deposition of new ECM by simultaneously stimulating the synthesis of new ECM, inhibiting the proteases that degrade matrix and modulating the numbers of integrin receptors to facilitate cell adhesion to the newly assembled matrix; 4) suppressing the proinflammatory effects of interleukin-1 and tumor necrosis factor; 5) regulating the action of platelet derived growth factor and fibroblast growth factor so that cell proliferation and angiogenesis are coordinated with matrix deposition; and 6) terminating the process when repair



is complete and the wound is closed (Border and Noble, *Scientific Amer. Sci. & Med.* 2:68-77 (1995)).

Interactions between RAS and TGF $\beta$  occur at both the systemic and molecular level. It has been shown that TGF $\beta$ 's action in causing ECM deposition in a healing wound is the same action that makes TGF $\beta$  a powerful fibrogenic cytokine. (Border and Noble, *New Engl. J. Med.* 331:1286-1292 (1994); and Border and Ruoslahti, *J. Clin. Invest.* 90:107 (1992)). Indeed, it is the failure to terminate the production of TGF $\beta$  that distinguishes normal tissue repair from fibrotic disease. RAS and TGF $\beta$  co-regulate each other's expression. Thus, both systems may remain active long after an emergency response has been terminated, which can lead to progressive fibrosis. The kidney is particularly susceptible to overexpression of TGF $\beta$ . The interrelationship of RAS and TGF $\beta$  may explain the susceptibility of the kidney to TGF $\beta$  overexpression and why pharmacologic suppression of RAS or inhibition of TGF $\beta$  are both therapeutic in fibrotic diseases of the kidney. (Noble and Border, *Sem. Nephrol.*, supra and Border and Noble, *Kidney Int.* 51:1388-1396 (1997)).

Activation of RAS and generation of angiotensin II (AII) are known to play a role in the pathogenesis of hypertension and renal and cardiac fibrosis. TGF $\beta$  has been shown to be a powerful fibrogenic cytokine, acting simultaneously to stimulate the synthesis of ECM, inhibit the action of proteases that degrade ECM and increasing the expression of cell surface integrins that interact with matrix components. Through these effects, TGF $\beta$  rapidly causes the deposition of excess ECM. AII infusion strongly stimulates the production and activation of TGF $\beta$  in the kidney. (Kagami et al., *J. Clin. Invest.* 93:2431-2437 (1994)). Angiotensin II also upregulates TGF $\beta$  production and increases activation when added to cultured vascular smooth muscle cells (Gibbons et al, *J. Clin. Invest.* 90:456-461 (1992)) and this increase is independent of pressure (Kagami et al., supra). AII also upregulates TGF $\beta$  receptors, even in the presence of exogenously added TGF $\beta$  which normally down-regulates its own receptors, leading to enhanced TGF $\beta$  signalling and enhanced fibronectin production (Kanai et al., *J. Am. Soc. Nephrol.* 8:518A (1997)). Blockade of AII reduces TGF $\beta$  overexpression in kidney and heart, and it is thought that TGF $\beta$  mediates renal and cardiac fibrosis associated with activation of RAS (Noble and Border, *Sem. Nephrol.* 17(5):455-466 (1997)), Peters et al., *Kidney International* 54 (1998)).

Blockade of AII using inhibitors of ACE slow the progression of renal fibrotic disease (see, e.g., Anderson et al., *J. Clin. Invest.* 76:612-619 (1985) and Noble and Border, *Sem. Nephrol.* 17(5):455-466 (1997)). What is not clear is whether angiotensin blockade reduces fibrosis solely through controlling glomerular hypertension and thereby glomerular injury, or whether pressure-independent as well as pressure-dependent mechanisms are operating. While ACE inhibitors and AII receptor antagonists have been shown to slow the progress of fibrotic diseases, they do not halt disease and TGF $\beta$  levels remain somewhat elevated. (Peters et al., supra).

Thus, RAS and TGF $\beta$  can be viewed as powerful effector molecules that interact to preserve systemic and tissue homeostasis. The response to an emergency such as tissue injury is that RAS and TGF $\beta$  become activated. Continued activation may result in chronic hypertension and progressive tissue fibrosis leading to organ failure. Because of the interplay between the RAS and TGF $\beta$ , and the effects of this interplay on tissue homeostasis, blockade of the RAS may be suboptimal to prevent or treat progressive fibrotic diseases such as diabetic nephropathy.

Components of the renin-angiotensin system act to further stimulate production of TGF $\beta$  and plasminogen activator inhibitor leading to rapid ECM accumulation. The protective effect of inhibition of the renin-angiotensin system in experimental and human kidney diseases correlates with the suppression of TGF $\beta$  production. (Noble and Border, *Sem. Nephrol.*, supra; and Peters et al., supra).

The renin molecule has been shown to enzymatically cleave angiotensinogen into Angiotensin I. The angiotensin I is then converted by Angiotensin Converting Enzyme ("ACE") to Angiotensin II which acts as an active metabolite and induces TGF $\beta$  production. Angiotensin II is an important modulator of systemic blood pressure. It has been thought that if you decrease hypertension by blocking AII's vasoconstrictor effects fibrotic disease is reduced.

In the glomerular endothelium, activation of RAS and TGF $\beta$  have been shown to play a role in the pathogenesis of glomerulonephritis and hypertensive injury. Volume (water) depletion and restriction of potassium have been shown to stimulate both production of renin and

5 TGF $\beta$  in the juxtaglomerular apparatus (JGA) of the kidney (Horikoshi et al., *J. Clin. Invest.* 88:2117-2122 (1992) and Ray et al., *Kidney Int.* 44:1006-1013 (1993)). Angiotensin blockade has also been shown to increase the production of renin. TGF $\beta$  has been shown to stimulate the release of renin from kidney cortical slices and cultured JG cells (Antonipillai et al., *Am. J. Physiol.* 265:F537-F541 (1993); Ray et al., *Contrib. Nephrol.* 118:238-248 (1996) and Veniant et al., *J. Clin. Invest.* 98:1996-19970 (1996)), suggesting that renin and TGF $\beta$  are coregulated. Other interactions between RAS and TGF $\beta$  include that AII induces the production of TGF $\beta$  in cultured cells and in vivo (Kagami et al., *supra*) and AII regulates expression of TGF $\beta$  receptors (Kanai et al., 1977, *supra*). It is thus likely that the fibrogenic effects that have been attributed to AII are actually mediated by TGF $\beta$ .

15 Another interplay between RAS and TGF $\beta$  is with the production of aldosterone. Aldosterone overproduction has been linked to hypertension and glomerulosclerosis. AII stimulates the production and release of aldosterone from the adrenal gland. In contrast, TGF $\beta$  suppresses aldosterone production and blocks the ability of AII to stimulate aldosterone by reducing the number of AII receptors expressed in the adrenal (Gupta et al., *Endocrinol.* 131:631-636 (1992)), and blocks the effects of aldosterone on sodium reabsorption in cultured renal collecting duct cells (Husted et al., *Am. J. Physiol. Renal, Fluid Electrolyte Physiol.* 267:F767-F775 (1994)). Aldosterone may have fibrogenic effects independent of AII, and may upregulate TGF $\beta$  expression. The mechanism of aldosterone's pathological effects is unknown but might be due to stimulation of TGF $\beta$  production in the kidney (Greene et al., *J. Clin. Invest.* 98:1063-1068 (1996)).

25 Prorenin or renin may have AII-independent actions to increase fibrotic disease. Prorenin overexpressing rats were found to be normotensive but to develop severe glomerulosclerosis (Veniant et al., *J. Clin. Invest.* 98:1996-19970 (1996)).

30 Human recombinant renin added to human mesangial cells induces marked upregulation of production of plasminogen activator inhibitors (e.g. PAI-1 and PAI-2) which block the generation of plasmin, a fibrinolytic enzyme important in the dissolution of clots after wounding generated from plasminogen by two enzymes called plasminogen activators, urokinase (u-PA)

and tissue plasminogen activator (t-PA). PAI-1 and 2 regulate U-PA and t-PA in turn. Plasmin appears to be a key mediator of extracellular matrix degradation, carrying out at least three functions important to extracellular matrix degradation. Plasmin directly degrades proteoglycan components of extracellular matrix, proteolytically activates metalloproteinases (MMPs) that, in turn, degrade collagens and other matrix proteins, and enzymatically inactivates tissue inhibitors of MMPs (TIMPs), releasing MMPs from inhibition of TIMPs, allowing them to proteolytically digest matrix proteins. (Baricos et al., *Kidney Int'l.* 47:1039-1047 (1995); Baricos et al., *J. Amer. Soc. Nephrol.* 10:790-795 (1999)). The net generation of active plasmin from the inactive precursor plasminogen results from a balance of the plasminogen activators and PAI-1 and 2, and other factors. PAI-1 binds to vitronectin. (Lawrence et al., *J. Biol. Chem.* 272:7676-7680 (1997)). Mutant PAI-1 molecules have been developed that have enhanced properties for PAI-1 binding to vitronectin molecules, but do not inhibit either t-PA or u-PA activity, resulting in an increase in the amount of the active form of plasmin. (See, WO 97/39028, Lawrence et al.). PAI-1 is increased in response to added TGF $\beta$  (Tomooka et al., *Kidney Int.* 42:1462-1469 (1992)).

It has been suggested that TGF $\beta$  enhances release of renin from storage granules in the juxtaglomerular apparatus of the kidney (Antonipillai et al., *Am. J. Physiol.* 265:F537-F541 (1993) and Ray et al., *Contrib. Nephrol.* 118:238-248 (1996)).

Thus, the interactions of RAS and TGF $\beta$  production form a complex system which impacts fibrotic ECM accumulation and the incidence of fibrotic disease. Various RAS components such as aldosterone, prorenin and renin may be connected with TGF $\beta$  production and fibrotic ECM accumulation. Any successful therapeutic regime must take into account these complex relationships to optimize inhibition of TGF $\beta$  to prevent and/or reduce ECM accumulation.

The multiple pathways resulting in TGF $\beta$  overexpression and fibrosis proposed from in vitro studies are depicted in Figure 1. (See, Kagami et al., *J. Clin. Invest.* 93:2431-2437 (1994); Gibbons et al., *J. Clin. Invest.* 90:456-461 (1992); Abboud, *Kidney Int.* 41:581-583 (1992); Ruiz-Ortega et al., *J. Am. Soc. Nephrol.* 5:683 (1994) abstract; Kim et al., *J. Biol. Chem.* 267:13702-

13707 (1992); Ohno et al., *J. Clin. Invest.* 95:1363-1369 (1995); Riser et al., *J. Clin. Invest.* 90:1932-1943 (1992); Riser et al., *J. Am. Soc. Nephrol.* 4:663 (1993); Ziyadeh et al., *J. Clin. Invest.* 93:536-542 (1994); Rocco et al., *Kidney Int.* 41:107-114 (1992); Flaumenhaft et al., *Advan. Pharmacol.* 24:51-76 (1993); Lopez-Armanda et al., *J. Am. Soc. Nephrol.* 5:812 (1994) abstract; Sahai et al., *J. Am. Soc. Nephrol.* 6:910 (1995); Remuzzi et al., *Kidney Int.* 1:2-15 (1997); and Remuzzi et al., *J. Am. Soc. Nephrol.* 9:1321-1332 (1998)). This diagram shows that a large number of factors implicated in kidney injury are believed to increase the production of TGF $\beta$ .

In fibrotic diseases overproduction of TGF $\beta$  results in excess accumulation of extracellular matrix which leads to tissue fibrosis and eventually organ failure. Accumulation of mesangial matrix is a histological indication of progressive glomerular diseases that lead to glomerulosclerosis and end-stage kidney disease (Klahr et al., *N. Engl. J. Med.* 318:1657-1666 (1988); Kashgarian and Sterzel, *Kidney Int.* 41:524-529 (1992)). Rats injected with antithymocyte serum are an accepted model of human glomerulonephritis and this model has demonstrated that overproduction of glomerular TGF $\beta$  can underlie the development of glomerulosclerosis (Okuda et al., *J. Clin. Invest.* 86:453-462 (1990); Border et al., *Nature (Lond.)* 346:371-374 (1990); Kagami et al., *Lab. Invest.* 69:68-76 (1993); and Isaka et al., *J. Clin. Invest.* 92:2597-2602 (1993)). Using cultured rat mesangial cells where the effects of Angiotensin II on glomerular pressure are not a factor, Angiotensin II has been shown to induce TGF $\beta$  production and secretion by mesangial cells, and this in turn has been shown to stimulate extracellular matrix production and deposition (Kagami et al., *J. Clin. Invest.* 93:2431-2437 (1994)). Increases in PAI-1 levels result in decreased degradation of extracellular matrix (Baricos et al., *Kidney Int.* 47:1039-1047 (1995)). Increases in TGF $\beta$  result in increased PAI-1 levels (Tomooka et al., *Kidney Int.* 42:1462-1469 (1992)). It has been demonstrated that decreasing TGF $\beta$  overexpression in a rat model of glomerulonephritis by in vivo injection of neutralizing antibodies to TGF $\beta$ , reduces TGF $\beta$  overexpression (Border et al., *Nature* 346:371-374 (1990)), and reduces PAI-1 deposition into the pathological matrix (Tomooka et al., *Kidney Int.* 42:1462-1469 (1992)). Therefore, decreases in TGF $\beta$  levels should result in decreased PAI-1 levels and increased degradation of extracellular matrix to ameliorate organ impairment and fibrotic disease. However, patients present with fibrotic disease that is well advanced in terms of build-up of

extra-cellular matrix (ECM). This is because abnormal organ function is undetectable until ECM accumulation is very advanced. For example, in the kidney, standard diagnostic tests do not provide an abnormal reading until about fifty percent of organ function has been lost.

5           The treatment of conditions associated with excess accumulation of ECM has also focused on decreasing stimuli to disease such as to lower blood pressure or, in the case of diabetic nephropathy to reduce plasma glucose levels. For example, current therapies for treating fibrotic disease in the kidney are limited to AII blockade using ACE inhibitors such as Enalapril or AII receptor antagonists such as Losartan. In addition, patients are encouraged to follow low protein diets since this regimen has some therapeutic value (Rosenberg et al., *J. Clin. Invest.* 10 85:1144-1149 (1992)). These therapies, at best, prolong organ function by only 1-2 years. This may be because of the multiple pathways that result in TGF $\beta$  overexpression or enhanced activity. Moreover, it is likely that current therapeutic strategies to reduce TGF $\beta$  overproduction may lead to upregulation of other pathways resulting in continued TGF $\beta$  overproduction. For example, when the action of AII is blocked, renin is upregulated which itself increases TGF $\beta$  production (see co-pending U.S. patent application, U.S. Serial No. 09/005,255, incorporated in its entirety herein). More recently, treatments aimed to halt the overproduction of TGF $\beta$  have been proposed (Border and Noble, *Kidney Internatl.* 54 (1998); and Peters et al., *Kidney Internatl.* 54 (1998)).

Therefore, the most promising therapeutic methods will need to increase ECM degradation to restore organ function as well as decrease TGF $\beta$  overproduction and/or activity. Enhanced degradation of excess accumulated ECM can be used to optimize overall reduction in levels of accumulated ECM to restore function to tissues and organs. Proteases that are able to degrade ECM are known. For example, the serine protease plasmin degrades ECM proteins and activates pro-metalloproteinases, in addition to degrading fibrin (Baricos et al., *supra*). One goal of therapeutic intervention to increase ECM degradation for treating fibrosis could be increasing plasmin in the region of excess ECM deposition.

30           There is a need for improved therapies to normalize TGF $\beta$  production, that take into account the multiple pathways that stimulate TGF $\beta$  production, to prevent or reduce excess

accumulation of ECM, to restore function to tissues and organs in which excess ECM has accumulated and/or to reduce scar formation at a wound site.

### SUMMARY OF THE INVENTION

Accordingly, the present invention provides methods for preventing or reducing the excess accumulation of extracellular matrix (ECM) associated with fibrotic conditions by inhibiting TGF $\beta$ , using a combination of agents that inhibit TGF $\beta$ , or by using a combination of agents to inhibit TGF $\beta$  and agents that cause the enhanced degradation of excess accumulated ECM.

The methods of the invention contemplate the use of agents that directly or indirectly inhibit TGF $\beta$  including direct inhibitors of TGF $\beta$  activity such as anti-TGF $\beta$  antibodies, proteoglycans such as decorin and ligands for TGF $\beta$  receptors, and/or indirect TGF $\beta$  inhibitors including aldosterone, inhibitors of aldosterone, inhibitors of angiotensin II, renin inhibitors, ACE inhibitors and AII receptor antagonists which act to decrease TGF $\beta$  production.

The methods of the invention also contemplate the use of agents that result in the enhanced degradation of excess accumulated matrix including proteases such as serine proteases including plasmin, metalloproteases, or protease combinations, and agents such as tPA, and PAI-1 mutants that increase the production and/or the activity of proteases such as plasmin.

The agents for use in the methods of the invention may be administered as inhibitory compounds in pharmaceutical formulations or as nucleic acid encoding the inhibitors delivered to suitable host cells. The nucleic acid may be directly introduced into a cell in vivo, for example into muscle tissue, or may be first introduced into a cell ex vivo to obtain a cell expressing the inhibitory agent or agents, and the cell then transplanted or grafted into a subject to inhibit or reduce excess accumulation of extracellular matrix.

The invention includes compositions for preventing or reducing the excess accumulation of ECM containing a combination of agents for inhibiting TGF $\beta$  or a combination of agents for inhibiting TGF $\beta$  and for degrading ECM.

5 **BRIEF DESCRIPTION OF THE DRAWINGS**

Figure 1 is a diagram depicting various pathways resulting in increased TGF $\beta$  production.

10 Figure 2 is a bar graph showing increases in TGF $\beta$  production by cultured human mesangial cells in response to renin, as described in Example I, infra.

Figure 3 is a bar graph showing the effect of blocking agents on TGF $\beta$ -production by human mesangial cells in response to renin, as described in Example II, infra.

15 Figure 4A and B are bar graphs showing dose dependent increases in TGF $\beta$  (Figure 4A) and Fn production (Figure 4B) with increases in HrRenin as described in Example IV, infra.

20 Figure 5A and B are bar graphs showing time courses of TGF $\beta$  (Figure 5A) and Fn production (Figure 5B) as described in Example IV, infra.

Figure 6A-C are bar graphs showing renin-induced increases in TGF $\beta$ , PAI-1 and Fn mRNAs over time as described in Example IV, infra.

25 Figure 7 is a bar graph showing the results of inhibitors that block renin's action to increase Angiotensin II, on the renin-induced increase in TGF $\beta$  production in adult human mesangial cells as described in Example IV, infra.

Figure 8A and B are photographs depicting the effects of tPA treatment on ECM accumulation in glomeruli as described in Example V, infra.

30 Figure 9A-D are bar graphs depicting the effects of tPA treatment on amounts of ECM



constituents (9A: FN EDA+; 9B:Laminin; 9C:Collagen I and 9D:Collagen IV) as determined by staining as described in Example V, infra.

Figure 10 is a bar graph showing the effects of tPA on glomerular mRNA expression at day 6 as described in Example V, infra.

Figure 11A and B are bar graphs showing the effects of tPA treatment on glomerular plasmin activity as described in Example V, infra.

Figure 12 is a bar graph demonstrating that injection of PAI-1 mutant results in increases in plasmin generation of nephritic glomeruli, as described in Example VII, infra.

Figure 13 is a bar graph demonstrating decreased accumulation of Collagen type I after administration of PAI-1 mutant, as described in Example VII, infra.

### DETAILED DESCRIPTION OF THE INVENTION

The present invention is based on the discovery that a combination of strategies may be warranted to prevent or treat conditions associated with the excess accumulation of extracellular matrix in tissues or organs, including fibrotic diseases and scarring resulting from TGF $\beta$  overproduction and/or activity. As previously reported, TGF $\beta$  overproduction may result from multiple pathways and require that more than one pathway be inhibited to achieve any clinically significant reduction in excess accumulation of extracellular matrix and amelioration of disease. For example, as disclosed in co-pending U.S. patent application, U.S. Serial No. 09/005,255, incorporated in its entirety herein, renin stimulates TGF $\beta$  production in cells capable of producing TGF $\beta$ , in an angiotensin-II and blood pressure-independent manner.

Optimal therapy of disorders associated with excess accumulation of ECM which causes organ impairment and ultimately failure, must take into account the multiple pathways of TGF $\beta$  production to effectively combat overproduction of TGF $\beta$ . Without such multifactorial strategy,

inhibition of one pathway of TGF $\beta$  production may be insufficient to block excess accumulation of extracellular matrix and can even result in an increase in the levels of TGF $\beta$  production by stimulation of one of the alternative pathways for its production.

5 While it is now known that multiple stimuli result in TGF $\beta$  overexpression and resulting excess accumulation of ECM, therapeutic strategies directly inhibiting TGF $\beta$ , such as the use of anti-TGF $\beta$  antibodies or TGF $\beta$  receptor antagonists, are being explored. However, because TGF $\beta$  has many beneficial actions such as immunosuppressive and immunomodulatory effects, as well as inhibition of epithelial cell growth which retards carcinogenesis (Markowitz, *Science* 10 268:1336-1338 (1995) and suppression of atherogenesis (Grainger et al., *Nature Med.* 1:74-79 (1995), these therapies may have unacceptable side-effects if administered at doses high enough to successfully stem fibrotic conditions. This has been shown in the TGF $\beta$ 1 null (knockout) mice which die of overwhelming inflammation at about 6 weeks of age (Letterio et al., *Science* 15 264:1936-1938 (1994); Kulkarni et al, *Proc. Natl. Acad. Sci. USA* 90:770-774 (1993) and Shull et al., *Nature* 359:693-699 (1992)), indicating that TGF $\beta$ 1 has significant beneficial roles in immune function. Multiple agents, inhibiting TGF $\beta$  directly, and/or inhibiting the disease-specific stimuli underlying TGF $\beta$  overexpression and/or activity, for example high glucose resulting from diabetes, may be required to adequately reduce TGF $\beta$ -associated excess accumulation of ECM, without causing harmful side-effects. Accordingly, it is a goal of the methods of the present invention to accomplish normalization of TGF production without harmful side effects and to prevent or reduce excess accumulation of ECM and ensuing fibrotic conditions. 20

In addition, degradation of accumulated ECM may be needed to restore tissue or organ 25 function that has been compromised by the presence of the excess accumulated ECM. Prevention or degradation of excess accumulated ECM can also prevent or reduce scar formation at the site of a wound.

The methods of the invention include using multiple agents to reduce the overproduction and/or activity of TGF $\beta$  and/or to block alternative pathways of TGF $\beta$  production to prevent or 30 reduce excess accumulation of ECM. The methods of the invention further include the use of

a combination of agents to reduce TGF $\beta$  overproduction and/or activity in combination with agents to enhance the degradation of excess, accumulated ECM. The methods are useful to prevent or reduce excess accumulation of extracellular matrix to ameliorate fibrotic conditions, and to restore or maintain normal tissue or organ function or skin appearance.

5

As used herein "excess accumulation of extracellular matrix" means the deposition of extracellular matrix components including, collagen, laminin, fibronectin and proteoglycans in tissue to an extent that results in impairment of tissue or organ function and ultimately, organ failure as a result of fibrotic disease. In addition, "excess accumulation of extracellular matrix" means the deposition of extracellular matrix components in the process commonly referred to as "scarring" or "scar formation," e.g. at a wound site. "Reducing the excess accumulation of extracellular matrix" means preventing excess accumulation of extracellular matrix, e.g. in tissue, organs or at a wound site, preventing further deposition of extracellular matrix and/or decreasing the amount of excess accumulated matrix already present, to maintain or restore tissue or organ function or appearance.

10

15

A variety of conditions are characterized by excess accumulation of extracellular matrix (collagen, fibronectin and other matrix components). Such conditions include, for example, but are not limited to, glomerulonephritis, adult or acute respiratory distress syndrome (ARDS), diabetes-associated pathologies such as diabetic kidney disease, fibrotic diseases of the liver, lung and post infarction cardiac fibrosis. Also included are fibrocystic diseases such as fibrosclerosis and fibrotic cancers such as cancers of the breast, uterus, pancreas or colon, and including fibroids, fibroma, fibroadenomas and fibrosarcomas.

20

25

There are also a number of medical conditions associated with an excess accumulation of extracellular matrix involving increased collagen, fibronectin and other matrix components. Such conditions include, for example, but are not limited to, post myocardial infarction, left ventricular hypertrophy, pulmonary fibrosis, liver cirrhosis, veno-occlusive disease, post-spinal cord injury, post-retinal and glaucoma surgery, post-angioplasty restenosis and renal interstitial fibrosis, arteriovenous graft failure, excessive scarring such as keloid scars and scars resulting from injury, burns or surgery.

30

As discussed, supra, it is known that TGF $\beta$  is indicated in the causation of fibrotic conditions. During normal tissue repair, TGF $\beta$  production is increased to stimulate the process of repair. When repair is complete, TGF $\beta$  production is reduced. If not reduced following normal tissue repair, the increased TGF $\beta$  overproduction can result in the development of excess extracellular matrix accumulation and fibrotic conditions. Thus, repeated tissue injury or a defect in TGF $\beta$  regulation leading to sustained TGF production results in excess accumulation of extracellular matrix.

As used herein "inhibition of TGF $\beta$ " includes inhibition of TGF $\beta$  activity, for example in causing excess deposition of ECM, as well as inhibition of TGF $\beta$  production resulting in overproduction and excess accumulation of ECM, regardless of the mechanism of TGF $\beta$  activity or overproduction. This inhibition can be caused directly, e.g. by binding to TGF $\beta$  or its receptors, for example by anti-TGF $\beta$  antibodies or TGF $\beta$  receptor antagonists, or can be caused indirectly, for example by inhibiting a pathway that results in TGF $\beta$  production, such as the renin pathway. Inhibition causes a reduction in the ECM producing activity of TGF $\beta$  regardless of the exact mechanism of inhibition.

As used herein a "TGF $\beta$  inhibitory agent" is an agent that directly or indirectly inhibits TGF $\beta$  binding to its receptors, such as a TGF $\beta$ -specific inhibitory agent, or an agent that blocks an alternative pathway of TGF $\beta$  production. The agent causes a reduction in the ECM producing activity of TGF $\beta$  regardless of the mechanism of its action. The agent can be nucleic acid encoding the TGF $\beta$  inhibitory agent such as a cDNA, genomic DNA, or an RNA or DNA encoding TGF $\beta$  inhibitory activity such as a TGF $\beta$  antisense RNA or DNA.

As used herein, a "TGF $\beta$ -specific inhibitory agent" means an agent containing TGF $\beta$  inhibiting activity, including agents that bind directly to TGF $\beta$  such as anti-TGF $\beta$  antibodies, or are a ligand for TGF $\beta$  which prevents it from binding to its receptors. A TGF $\beta$ -specific inhibiting agent also includes a nucleic acid encoding a particular TGF $\beta$ -specific inhibitory agent such as a cDNA, genomic DNA or an RNA or DNA encoding TGF $\beta$ -specific inhibitory activity such as a TGF $\beta$  antisense RNA or DNA.

Agents that bind directly to TGF $\beta$  are known and include anti-TGF $\beta$  antibodies such as anti-TGF $\beta$ 1 antibodies (Genzyme, Cambridge, MA) and antibodies which bind both TGF $\beta$ 1 and TGF $\beta$ 2 (Dasch et al., U.S. Patent No. 5,571,714), proteoglycans such as decorin, biglycan and fibromodulin, and the nucleic acids encoding such agents.

Antibodies to inhibit TGF $\beta$ , renin or other molecules, for use in the present invention, can be prepared according to methods well established in the art, for example by immunization of suitable host animals with the selected antigen, e.g. TGF $\beta$ . For descriptions of techniques for obtaining monoclonal antibodies see, e.g. the hybridoma technique of Kohler and Milstein (*Nature* 256:495-497 (1975)), the human B-cell hybridoma technique (Kosbor et al., *Immunol. Today* 4:72 (1983); Cole et al., *Proc. Nat'l. Acad. Sci. USA*, 80:2026-2030 (1983)) and the EBV-hybridoma technique (Cole et al., *Monoclonal antibodies and Cancer Therapy*, Alan R. Liss, Inc., pp. 77096 (1985)). Such antibodies may be of any immunoglobulin class including IgG, IgM, IgE, IgA, IgD and any subclass thereof. The hybridoma producing the monoclonal antibody may be cultivated *in vitro* or *in vivo*. Suitable host animals include, but are not limited to, rabbits, mice, rats, and goats. Various adjuvants may be used to increase the immunological response to the host animal, depending on the host species, including, but not limited to, Freund's (complete and incomplete), mineral gels such as aluminum hydroxide, surface active substances such as pluronic polyols, polyanions, peptides, oil emulsions, keyhole limpit, hemocyanin, dinitrophenol and potentially useful human adjuvants such as BCG (*Bacille Calmette-Guerin*) and *Cornebacterium parvum*. Antibodies as used herein includes non-human, chimeric (different species), humanized (see Borrebaeck, *Antibody Engineering: A Practical Guide*, W.H. Freeman and Co., New York, 1991), human and single-chain antibodies, as well as antibody fragments including but not limited to the F(ab')<sub>2</sub> fragments that can be produced by pepsin digestion of antibody molecules and Fab fragments that can be generated by reducing disulfide bridges of the F(ab')<sub>2</sub> fragments. Alternatively, Fab expression libraries may be constructed (*Science* 246:1275-1281 (1989)) to permit the rapid and easy identification of monoclonal Fab fragments having the desired specificity.

An indirect TGF $\beta$  inhibitor would inhibit the synthesis or secretion of TGF $\beta$  or sequester it away from its target cells. Such inhibitors include, but are not limited to, inhibitors of

Angiotensin Converting Enzyme ("ACE"), antagonists of the AII receptor such as Losartan<sup>™</sup> and Cozar<sup>™</sup> (Merck), and aldosterone inhibitors such as Spironolactone<sup>™</sup> (Sigma Chemical Co., St. Louis, Mo, Product # S 3378) that would otherwise result in increased TGF $\beta$  production.

5 Also included within the scope of TGF $\beta$  inhibitors of the invention are nucleic acids that include antisense oligonucleotides that block the expression of specific genes within cells by binding a complementary messenger RNA (mRNA) and preventing its translation (See review by Wagner, *Nature* 372:332-335 (1994); and Crooke and Lebleu, *Antisense Research and Applications*, CRC Press, Boca Raton (1993)). Gene inhibition may be measured by determining  
10 the degradation of the target RNA. Antisense DNA and RNA can be prepared by methods known in the art for synthesis of RNA including chemical synthesis such as solid phase phosphoramidite chemical synthesis or in vitro and in vivo transcription of DNA sequences encoding the antisense RNA molecule. The DNA sequences may be incorporated into vectors with RNA polymerase promoters such as the T7 or SP6 polymerase promoters. Alternatively, antisense cDNA constructs that synthesize antisense RNA constitutively or inducibly can be introduced into cell lines. The potency of antisense oligonucleotides for inhibiting TGF $\beta$  may be enhanced using various methods including 1) addition of polylysine (Leonetti et al., *Bioconj. Biochem.* 1:149-153 (1990)); 2) encapsulation into antibody targeted liposomes (Leonetti et al., *Proc. Natl. Acad. Sci. USA* 87:2448-2451 (1990) and Zelphati et al., *Antisense Research and Development* 3:323-338 (1993)); 3) nanoparticles (Rajaonarivony et al., *J. Pharmaceutical Sciences* 82:912-917 (1993) and Haensler and Szoka, *Bioconj. Chem.* 4:372-379 (1993)), 4) the use of cationic acid liposomes (Felgner et al., *Proc. Natl. Acad. Sci. USA* 84:7413-7417 (1987); Capaccioli et al., *Biochem. Biophys. Res. Commun.* 197:818-825 (1993); Boutorine and Kostina, *Biochimie* 75:35-41 (1993); Zhu et al., *Science* 261:209-211 (1993); Bennett et al., *Molec. Pharmac.* 41:1023-1033 (1992) and Wagner, *Science* 280:1510-1513 (1993)); and 5) Sendai virus derived liposomes (Compagnon et al., *Exper. Cell Res.* 200:333-338 (1992) and Morishita et al., *Proc. Natl. Acad. Sci. USA* 90:8474-8478 (1993)), to deliver the oligonucleotides into cells. Recent techniques for enhancing delivery include the conjugation of the antisense oligonucleotides to a fusogenic peptide, e.g. derived from an influenza hemagglutinin envelop protein (Bongartz et al., *Nucleic Acids Res.* 22(22):4681-4688 (1994)).  
25  
30

Additional suitable TGF $\beta$  inhibitory agents can be readily obtained using methods known in the art to screen candidate agent molecules for binding to TGF $\beta$ , such as assays for detecting the ability of a candidate agent to block binding of radiolabeled human TGF $\beta$  to cells such as human mesangial cells. Alternatively, candidate compounds may be tested for the ability to inhibit TGF $\beta$  production by mesangial cells using an enzyme-linked immunosorbent assay (ELISA), for example using the R & D Systems (Minneapolis, MN) TGF $\beta$  ELISA assay kit (Cat. No. DB 100) (for methods see, e.g. Uotila et al., *J. Immunol. Methods* 42:11 (1981)).

Suitable TGF $\beta$ -specific inhibitory agents can also be developed by known drug design methods, e.g. using structural analysis of the TGF $\beta$  molecule employing methods established in the art, for example, using X-ray crystallography to analyze the structure of the complex formed by TGF $\beta$  and one of its known inhibitors (see, e.g. Sielecki et al., *supra*; Rahuel et al., *supra*, Badasso et al., *supra* and Dhanaraj et al., *supra*), and/or by modifying known TGF $\beta$  antagonists i.e. "lead compounds," to obtain more potent inhibitors and compounds for different modes of administration (i.e. oral vs. intravenous) (see, e.g. Wexler et al., *Amer. J. Hyper.* 5:209S-220S (1992)-development of AII receptor antagonists from Losartan<sup>™</sup>). For such procedures large quantities of TGF $\beta$  can be generated using recombinant technology or purchased commercially (R & D Systems).

In addition to TGF $\beta$  inhibitory agents, agents that result in the degradation of ECM are contemplated for use in the invention. Such agents include serine proteases such as plasmin and metalloproteinases, and protease combinations such as Wobenzym (Mucos Pharma, Geretsried, Germany). In addition, the present inventors have discovered that agents such as tPA can be used to increase the amount of active proteases in vivo to increase degradation of ECM accumulated in organs and tissues. Tissue plasmin activator (tPA, Activase, Genentech, S. San Francisco, CA) has been shown to dissolve clots associated with myocardial infarction and stroke. The present inventors theorized that tPA might be helpful in increasing plasmin to reduce accumulated ECM. Shown herein is the use of recombinant tPA (rtPA) to increase the generation of plasmin in vivo to degrade ECM (Example V, *infra*).

In addition, new proteases or agonists of protease production and/or activity may be discovered or developed using rational drug design and used to degrade ECM according to the methods of the present invention.

5 The present inventors have also discovered that PAI mutants, such as the PAI-1 mutants disclosed in WO 97/39028 by Lawrence et al., incorporated by reference in its entirety herein, may be used to increase the amount of active plasmin to enhance degradation of ECM accumulated in organs and tissues. These PAI-1 mutants fail to inhibit plasminogen activators, yet retain significant vitronectin binding affinity. Additional PAI-1 mutants for use in the  
10 methods of the invention may be obtained and tested for the ability to bind vitronectin while failing to inhibit plasminogen activators (Lawrence et al., *J. Biol. Chem.* 272:7676-7680 (1997)). PAI-1 binding to vitronectin may be determined either functionally (Lawrence et al., *J. Biol. Chem.* 265:20293-20301 (1990)) or in a vitronectin specific ELISA (Lawrence et al., *J. Biol. Chem.* 269:15223-15228 (1994)). The ability of PAI-1 to inhibit plasminogen activators may be evaluated using chromogenic assays as described by Sherman et al., *J. Biol. Chem.* 270:9301-9306 (1995)).

15 In the methods of the invention, the TGF $\beta$  inhibitory agents are administered concurrently or sequentially. For example, an anti-TGF $\beta$  antibody is administered with an anti-renin agent. The inhibitory agents will localize at sites of TGF $\beta$  overproduction, e.g. organs such as the kidneys. The inhibitory agents may be labelled, using using known radiolabelling methods to detect their localization in a subject after administration. The agents may also be conjugated to targeting molecules such as antibodies to ECM components to improve localization of the agents after administration to the sites of TGF $\beta$  overproduction and/or excess accumulation of ECM in  
20 a subject.

25 In another embodiment of the methods of the invention, TGF $\beta$  inhibitory agents are administered concurrently or sequentially with at least one agent that degrades accumulated ECM, for example, a serine protease such as plasmin. Alternatively, an agent that induces protease production, such as tPA, is administered to increase protease production at the site(s)  
30 of accumulated ECM. tPA binds fibrin (Rondeau et al., *Clinical Nephrol.* 33:55-60 (1990)) and



thus will localize in fibrotic areas where the increased protease production is desired.

5 In one embodiment of the invention, at least one TGF $\beta$ -inhibitory agent is administered to a subject having existing excess accumulation of ECM in tissues or organs, or at high risk for such accumulation to reduce or prevent excess accumulation of ECM. For example, individuals at risk for developing fibrotic conditions, such as a person having or at high risk for diabetes, high blood pressure, autoimmune disease (e.g. lupus) and inflammatory diseases, can be scanned using known medical procedures including tissue biopsies of kidney, lung or liver, to determine whether ECM has accumulated in these organs. If the agent is TGF $\beta$ -specific, it binds to circulating TGF $\beta$  or tissue TGF $\beta$ . If the agent indirectly inhibits TGF $\beta$ , for example an anti-renin agent, it reduces the amount of TGF $\beta$  produced. As a result of the administration of agents that directly or indirectly inhibits TGF $\beta$ , ECM that has accumulated at the time of diagnosis or treatment, as well as further accumulation of ECM is reduced. Moreover, in high risk individuals the methods of the invention for inhibiting TGF $\beta$  overproduction with multiple agents can result in prevention of excess accumulation of ECM and the development of fibrotic conditions.

10 In another embodiment of the methods of the invention, at least one TGF $\beta$  inhibitory agent is administered to a subject having an existing excess accumulation of ECM in tissues or organs together with at least one agent to degrade accumulated ECM. The ECM degradation is accomplished using a protease, or an agent that enhances production or the activity of ECM degrading agents such as proteases. As a result of the administration of these agents, excess matrix accumulated at the time of diagnosis or treatment, as well as further excess accumulation of ECM is reduced.

20 In addition to the use of molecules such as antibodies and purified compounds such as decorin, nucleic acid encoding the TGF $\beta$  inhibitory agents and nucleic acid encoding the agent to directly or indirectly degrade accumulated ECM, are administered to the subject to permit the agents to be expressed and secreted, for inhibiting TGF $\beta$  and degrading accumulated ECM. The nucleic acid may be introduced into cells in the subject, for example using a suitable delivery vehicle such as an expression vector or encapsulation unit such as a liposome, or may be introduced directly through the skin, for example in a DNA vaccine.

Alternatively, the nucleic acids encoding the agents are introduced into a cell ex vivo and the cells expressing the nucleic acids are introduced into a subject, e.g. by implantation procedures, to deliver the agents in vivo. Multiple agents can be introduced into a delivery vehicle or in separate vehicles.

### Gene Therapy Methods

Methods for obtaining nucleic acids encoding TGF $\beta$  inhibitory agents and ECM degrading agents are known in the art. Following is a general description of methods of using the nucleic acids in gene therapy to reduce excess accumulation of ECM.

In one embodiment of the invention, gene therapy is contemplated using nucleic acids encoding the TGF $\beta$  inhibitory agents and/or the ECM degradation agent, introduced into cells in a subject to suppress TGF $\beta$  overproduction and to degrade accumulated ECM. Gene transfer into cells of these nucleic acids is contemplated in the methods of the invention.

### Nucleic Acids

Large amounts of the nucleic acid sequences encoding the TGF $\beta$ -inhibiting agents and/or the ECM degradation agents may be obtained using well-established procedures for molecular cloning and replication of the vector or plasmid carrying the sequences in a suitable host cell. DNA sequences encoding a specific agent can be assembled from cDNA fragments and oligonucleotide linkers, or from a series of oligonucleotides to provide a synthetic inhibitor agent gene and/or ECM degradation gene which can be expressed. Such sequences are preferably provided in an open reading frame uninterrupted by internal non-translated sequences or introns, which are typically present in eukaryotic genes. Genomic DNA containing the relevant sequences can also be used. Sequences of non-translated DNA may be present 5' to 3' from the open reading frame, where such sequences do not interfere with manipulation or expression of the coding regions. Either complete gene sequences or partial sequences encoding the desired agents are employed.

5 The nucleic acid sequences encoding the agents can also be produced in part or in total by chemical synthesis, e.g. by the phosphoramidite method described by Beaucage and Carruthers, *Tetra Letts.* 22:1859-1862 (1981) or the triester method (Matteucci et al., *J. Am. Chem. Soc.* 103:3185 (1981) and may be performed on commercial automated oligonucleotide synthesizers. A double-stranded fragment may be obtained from the single-stranded product of chemical synthesis either by synthesizing the complementary strand and annealing the strand together under appropriate conditions, or by synthesizing the complementary strand using DNA polymerase with an appropriate primer sequence.

## 10 Gene Transfer

15 For gene transfer, the key steps are 1) to select the mode of delivery, e.g. a proper vector for delivery of the inhibitor genes to the subject, 2) administer the nucleic acid to the subject; and 3) achieve appropriate expression of the transferred gene for satisfactory durations. Methods for gene transfer are known in the art. The methods described below are merely for purposes of illustration and are typical of those that can be used to practice the invention. However, other procedures may also be employed, as is understood in the art. Most of the techniques to construct delivery vehicles such as vectors and the like are widely practiced in the art, and most practitioners are familiar with the standard resource materials which describe specific conditions, reagents and procedures. The following paragraphs may serve as a guideline.

25 Techniques for nucleic acid manipulation are well known. (See, e.g. *Annual Rev. of Biochem.* 61:131-156 (1992)). Reagents useful in applying such techniques, such as restriction enzymes and the like, are widely known in the art and commercially available from a number of vendors.

30 The natural or synthetic nucleic acid coding for the inhibitors for expression in a subject may be incorporated into vectors capable of introduction into and replication in the subject. In general, nucleic acid encoding the selected inhibitor molecules and/or ECM degradation molecules are inserted using standard recombinant techniques into a vector containing appropriate transcription and translation control sequences, including initiation sequences

operably linked to the gene sequence to result in expression of the recombinant genes in the recipient host cells. "Operably linked" means that the components are in a physical and functional relationship permitting them to function in their intended manner.

5 For example, a promoter is operably linked to a coding sequence if the promoter effects its transcription or expression.

Sequences encoding selected inhibitor and/or degradation genes will include at least a portion of the coding sequence sufficient to provide the TGF $\beta$  inhibitory or ECM degradation activity in the expressed molecule. For example, in the case of a renin inhibitor, a portion of the coding sequence that enables the inhibitor to bind to renin can be used. Methods for determining such portions or "domains" including binding domains of molecules, are known in the art (See, e.g., Linsley et al., *Proc. Natl. Acad. Sci. USA* 87:5031-5035 (1990)). It is possible that it may be necessary to block both the renin enzymatic site and the renin-cell binding domain in order to effectively prevent the stimulus to TGF $\beta$  overproduction by renin. In such case, renin antisense molecules can be prepared using standard methods to accomplish complete blockade.

The selected nucleic acid sequences are inserted into a single vector or separate vectors. More than one gene encoding a selected agent, or portion thereof containing the desired activity, may be inserted into a single vector or into separate vectors for introduction into the host cells. Alternatively, these sequences can be administered as naked nucleic acid sequences or as part of a complex with other molecules, e.g. liposomes.

A variety of expression vectors and gene transfer methods useful for obtaining expression of selected molecule in recipient cells are well known in the art, and can be constructed using standard ligation and restriction techniques (see, for example, Sambrook et al., *Molecular Cloning: A Laboratory Manual*, Cold Spring Harbor Laboratory Press, 1989; Maniatis et al., *Molecular Cloning: A Laboratory Manual*, Cold Spring Harbor, New York (1982), Kriegler, *Gene Transfer and Expression: A Laboratory Manual* (W.H. Freeman and Co., New York, NY 1990) and Wu, *Methods in Enzymol.* (Academic Press, New York, NY 1993), each of which is incorporated by reference herein). The choice of vector or method depends on several factors

such as the particular molecule to be expressed.

Suitable vectors may be plasmid or viral vectors (Kaufman, in *Gene Expression Technology*, Goeddel (Ed.) (1991)) including baculoviruses, adenoviruses, poxviruses (Moss, *Current Opin. Biotech.* 3:518-522 (1993)), retrotransposon vectors (Cook et al., *Bio/Technology* 9:748-751 (1991) and Chakraborty et al., *FASEB J.* 7:971-977 (1993)) adeno-associated viruses (AAV) (Yei et al., *Gene Therapy* 1:192-200 (1994) and Smith et al., *Nat. Genet.* 5:397-402 (1993)), herpes virus and retrovirus vectors (Price et al., *Proc. Natl. Acad. Sci. USA* 84:156-160 (1987); Naviaux and Verma, *Current Opinion in Biotechnol.* 3:540-547 (1992); Hodgson and Chakraborty, *Curr. Opin. Thera. Patients* 3:223-235 (1993)) such as the MMLV based replication incompetent vector pMV-7 (Kirschmeier et al., *DNA* 7:219-225 (1988)), as well as human and yeast artificial chromosomes (HACs and YACs) (Huxley, *Gene Therapy* 1:7-12 (1994) and Huxley et al., *Bio/Technology* 12:586-590 (1994)). Plasmid expression vectors include plasmids including pBR322, pUC or Bluescript<sup>™</sup> (Stratagene, San Diego, CA).

Vectors containing the nucleic acid encoding the selected agents are preferably recombinant expression vectors in which high levels of gene expression may occur, and which contain appropriate regulatory sequences for transcription and translation of the inserted nucleic acid sequence. Regulatory sequences refer to those sequences normally associated (e.g. within 50 kb) of the coding region of a locus which affect the expression of the gene (including transcription, translation, splicing, stability or the like, of the messenger RNA). A transcriptional regulatory region encompasses all the elements necessary for transcription, including the promoter sequence, enhancer sequence and transcription factor binding sites. Regulatory sequences also include, inter alia, splice sites and polyadenylation sites. An internal ribosome entry site (IRES) sequence may be placed between recombinant coding sequences to permit expression of more than one coding sequence with a single promoter.

Transcriptional control regions include: the SV40 early promoter region, the cytomegalovirus (CMV) promoter (human CMV IE94 promoter region (Boshart et al., *Cell* 41:521-530 (1985)); the promoter contained in the 3' long terminal repeat of Rous Sarcoma Virus or other retroviruses; the herpes thymidine kinase promoter; the regulatory sequences of the

methallothionein gene; regions from the human IL-2 gene (Fujita et al., *Cell* 46:401-407 (1986)); regions from the human IFN gene (Ciccarone et al., *J. Immunol.* 144:725-730 (1990)); regions from the human IFN gene (Shoemaker et al., *Proc. Natl. Acad. Sci. USA* 87:9650-9654 (1990)); regions from the human IL-4 gene (Arai et al., *J. Immunol.* 142:274-282 (1989)); regions from the human lymphotoxin gene (Nedwin et al., *Nucl. Acids. Res.* 13:6361-6373 (1985)); regions from the human granulocyte-macrophage CSF gene (GM-CSF) (Miyatake et al., *EMBO J.* 4:2561-2568 (1985)) and others. When viral vectors are used, recombinant coding sequences may be positioned in the vector so that their expression is regulated by regulatory sequences such as promoters naturally residing in the viral vector.

Operational elements for obtaining expression may include leader sequences, termination codons and other sequences needed or preferred for the appropriate transcription and translation of the inserted nucleic acid sequences. Secretion signals may also be included whether from the native inhibitor or from other secreted polypeptides, which permit the molecule to enter cell membranes and attain a functional conformation. It will be understood by one skilled in the art that the correction type and combination of expression control elements depends on the recipient host cells chosen to express the molecules ex vivo. The expression vector should contain additional elements needed for the transfer and subsequent replication of the expression vector containing the inserted nucleic acid sequences in the host cells. Examples of such elements include, but are not limited to, origins of replication and selectable markers. Additionally, elements such as enhancer sequences, for example CMV enhancer sequences, may be used to increase the level of therapeutic gene expression (Armstrong. *Proc. Natl. Acad. Sci. USA* 70:2702 (1973)).

The vector may contain at least one positive marker that enables the selection of cells carrying the inserted nucleic acids. The selectable molecule may be a gene which, upon introduction into the host cell, expresses a dominant phenotype permitting positive selection of cells carrying the gene ex vivo. Genes of this type are known in the art and include, for example, drug resistance genes such as hygromycin-B phosphotransferase (hph) which confers resistance to the antibiotic G418; the aminoglycoside phosphotransferase gene (*neo* or *aph*) from Tn5 which codes for resistance to the antibiotic G418; the dihydrofolate reductase (DHFR) gene; the

adenosine deaminase gene (ADA) and the multi-drug resistance (MDR) gene.

Recombinant viral vectors are introduced into host cells using standard techniques. Infection techniques have been developed which use recombinant infectious virus particles for gene delivery into cells. Viral vectors used in this way include vectors derived from simian virus 40 (SV40; Karlsson et al., *Proc. Natl. Acad. Sci. USA* 82:158 (1985)); adenoviruses (Karlsson et al., *EMBO J.* 5:2377 (1986)); vaccinia virus (Moss et al., *Vaccine* 6:161-3 (1988)); and retroviruses (Coffin, in Weiss et al. (Eds.), *RNA Tumor Viruses*, 2nd Ed., Vol. 2, Cold Spring Laboratory, NY, pp. 17-71 (1985)).

Nonreplicating viral vectors can be produced in packaging cell lines which produce virus particles which are infectious but replication defective, rendering them useful vectors for introduction of nucleic acid into a cell lacking complementary genetic information enabling encapsidation (Mann et al., *Cell* 33:153 (1983); Miller and Buttimore, *Mol. Cell. Biol.* 6:2895 (PA317, ATCC CRL9078). Packaging cell lines which contain amphotrophic packaging genes able to transduce cells of human and other species origin are preferred.

Vectors containing the inserted inhibitor genes or coding sequences are introduced into host cell using standard methods of transfection including electroporation, liposomal preparations, Ca-PH-DNA gels, DEAE-dextran, nucleic acid particle "guns" and other suitable methods.

In additional to various vectors including viral vectors, other delivery systems may be used including, but not limited to, microinjection (DePamphilis et al., *BioTechnique* 6:662-680 (1988)); liposomal mediated transfection (Felgner et al., *Proc. Natl. Acad. Sci. USA* 84:7413-7417 (1987); Felgner and Holm, *Focus* 11:21-25 (1989) and Felgner et al., *Proc. West. Pharmacol. Soc.* 32:115-121 (1989)); use of naked or particle mediated DNA transfer and other methods known in the art. Recently, cationic liposomes have been used to enhance transfection (Felgner et al., *Nature* 349:351 (1991); Zhu et al., *Science* 261:209 (1993)).

Suitable host cells for gene transfer consist of vertebrate cells such as fibroblasts, keratinocytes, muscle cells, mesangial cells (see, Kitamura et al., *Kidney Int.* 48:1747-1757 (1995)), and any other suitable host cell including so-called universal host cells, i.e. cells obtained from a different donor than the recipient subject but genetically modified to inhibit rejection by the subject. Autologous cells are preferred, but heterologous cells are encompassed within the scope of the invention.

Expression of the selected TGF $\beta$  inhibitor genes after introduction into the host cells is confirmed using standard methods. For example, expression of TGF $\beta$ -specific inhibitory agents can be determined by assaying for the ability of the supernatant from transfected cells to inhibit the binding of radiolabeled TGF $\beta$  to human mesangial cells using Fluorescent Activated Cell Sorting (FACS) or ELISA. Expression from host cells of an agent that inhibits TGF $\beta$  indirectly, such as Losartan, can be confirmed by detecting a decrease in fibronectin production by mesangial cells exposed to supernatant from transfected cells, relative to controls. Expression of genes encoding ECM degrading agents can be determined using, for example, an in vitro system using mesangial cells cultured on a ECM substrate such as Matrigel<sup>™</sup> (Collaborative Research, Inc., Bedford, MA) that contains the major components of the mesangial matrix, including laminin, type IV collagen, entactin and heparan sulfate proteoglycan, as described by Baricos et al., *Kidney Internatl.* 47:1039-1047 (1995)). The ECM substrate is radiolabeled, and ECM degradation by the product of an expressed gene from transfected host cells is determined by measuring the release of radioactivity from the ECM into serum-free medium. These assay systems may also be employed to screen candidate TGF $\beta$  inhibiting and ECM degrading agents.

#### Administration of TGF $\beta$ Inhibitory Agents and Agents Degrading Accumulated ECM

Agents for inhibiting TGF $\beta$  and agents for degrading accumulated ECM are suspended in physiologically compatible pharmaceutical carriers, such as physiological saline, phosphate-buffered saline, or the like to form physiologically acceptable aqueous pharmaceutical compositions for administration to a subject. Parenteral vehicles include sodium chloride solution, Ringer's dextrose, dextrose and sodium chloride and lactated Ringer's solution. Other substances may be added as desired, such as antimicrobials.



The TGF $\beta$  inhibiting and ECM degrading agents may be administered together or apart, simultaneously or sequentially, to carry out the methods of the invention.

Modes of administration of the TGF $\beta$  inhibitory agents and ECM degrading agents are those known in the art for therapeutic agents and include parenteral, for example, intravenous (e.g. for antibody inhibitors or proteases), intraperitoneal, intramuscular, intradermal, and epidermal including subcutaneous and intradermal, oral (e.g. small molecule renin and TGF $\beta$  antagonists), or applied to mucosal surfaces, e.g. by intranasal administration using inhalation of aerosol suspensions, and by implanting to muscle or other tissue in the subject (e.g. for gene transfer of nucleic acid expressing renin and/or TGF $\beta$  inhibitors). Suppositories and topical preparations are also contemplated.

The TGF $\beta$  inhibitory and ECM degrading agents are introduced in amounts sufficient to prevent or reduce excess accumulation of extracellular matrix in susceptible tissues and organs including, but not limited to, lung and kidney tissue. Before or after administration, if necessary to prevent or inhibit the subject's immune response to the vehicles carrying the inhibitors, immunosuppressant agents may be used. Alternatively, the vehicles carrying the TGF $\beta$  inhibitory and ECM degrading agents can be encapsulated.

The most effective mode of administration and dosage regimen for the TGF $\beta$  inhibitory and ECM degrading agents for use in the methods of the present invention depend on the extent of TGF $\beta$  overproduction, the severity of the accumulation of extracellular matrix and resulting impairment of tissue or organ function, the subject's health, previous medical history, age, weight, height, sex and response to treatment and the judgment of the treating physician. Therefore, the amount of TGF $\beta$  inhibitory and ECM degrading agents to be administered, as well as the number and timing of subsequent administrations, are determined by a medical professional conducting therapy based on the response of the individual subject. Initially, such parameters are readily determined by skilled practitioners using appropriate testing in animal models for safety and efficacy, and in human subjects during clinical trials of candidate therapeutic formulations. Suitable animal models of human fibrotic conditions are known (see, e.g. Border and Noble, *New Eng. J. Med.* 331:1286-1292 (1994), incorporated by reference

herein).

After administration, the efficacy of the therapy using the methods of the invention is assessed by various methods including biopsy of kidney, lung or liver or other tissue to detect the amount of extracellular matrix accumulated. An absence of significant excess accumulation of ECM, or a decrease in the amount or expansion of ECM in the tissue or organ will indicate the desired therapeutic response in the subject. Preferably, a non-invasive procedure is used to detect a therapeutic response. For example, changes in TGF $\beta$  activity can be measured in plasma samples taken before and after treatment with an inhibitor (see, Eltayeb et al., *J. Am. Soc. Nephrol.* 8:110A (1997)), and biopsy tissue can be used to individually isolate diseased glomeruli which are then used for RNA isolation. mRNA transcripts for TGF $\beta$ , and extracellular matrix components (e.g. collagen) are then determined using reverse transcriptase-polymerase chain reaction (RT-PCR) (Peten et al., *J. Exp. Med.* 176:1571-1576 (1992)).

#### Advantages of the Invention

The invention provides improved treatment and prevention of fibrotic conditions associated with overproduction of TGF $\beta$  and excess accumulation of ECM in tissues and/or organs resulting in impaired function, or scarring, by reducing TGF $\beta$  overproduction directly and that resulting from multiple biological pathways, to effectively inhibit the TGF $\beta$  induced component of extracellular matrix deposition, and by increased degradation of ECM using degrading agents.

The therapeutic effects of the invention result from a reduction in or prevention of the TGF $\beta$ -induced excess accumulation of extracellular matrix in tissues and/or organs, and when combined with ECM degrading agents, from the increased degradation of ECM over time.

The following examples are presented to demonstrate the methods of the present invention and to assist one of ordinary skill in using the same. The examples are not intended in any way to otherwise limit the scope of the disclosure of the protection granted by Letters Patent granted hereon.

### EXAMPLE I

#### DEMONSTRATION THAT RENIN UPREGULATES TGF $\beta$ IN HUMAN MESANGIAL CELLS

Normal fetal human mesangial cells (Clonetics Corp., Clonetics, Walkersville, MD) passaged 5 to 8 times, were plated (3,000 cell/cm<sup>2</sup>) in 12 well plates in 2ml of medium (Mesangial Basal Medium (Clonetics Corp.) containing 5% FCS, 10  $\mu$ g/ml penicillin and 100  $\mu$ g/ml streptomycin) and allowed to grow to confluence for 48 hours at 37°C, 5% CO<sub>2</sub>. Cultures were washed three times using sterile phosphate buffered saline at room temperature and then 2 ml/well of serum free MBM medium to induce quiescence. After 48 hours, the serum-free medium was removed and 2 ml/well of fresh serum-free medium was added. Human recombinant renin (Hoffman-La Roche Ltd., Basel, Switzerland) in concentrations from 10<sup>-6</sup> to 10<sup>-12</sup> M was added to each well. A blank and 5 ng/ml of TGF $\beta$  (R & D Systems, Minneapolis, MN) were used as controls. Cells and supernatants were harvested by centrifugation after 24 hrs of culture and frozen at -70°C until analysis. The total production and release of TGF $\beta$  into the culture supernatant was measured using an ELISA kit (R & D Systems). Induction of PAI-1 and fibronectin in the supernatant are also measured using anti-PAI-1 and anti-fibronectin antibodies in an ELISA to provide further confirmation of the inhibition of TGF $\beta$ . TGF $\beta$ , fibronectin and PAI-1 mRNA are measured using semi-quantitative RT-PCR.

##### (1) Determination of Dose Dependency of Renin Induction of TGF $\beta$

As shown in Figure 2, renin increases the TGF $\beta$  production by cultured human mesangial cells in a dose-dependent manner.

## EXAMPLE II

### DEMONSTRATION OF THE EFFECT OF INHIBITING RENIN ON TGF $\beta$ PRODUCTION BY HUMAN MESANGIAL CELLS

Renin inhibitor Ro42-5892 (Hoffman-LaRoche, Basel, Switzerland), Losartan<sup>™</sup> (Merck Pharmaceuticals, West Point, PA), Enalapril<sup>™</sup> (Sigma Chemical Co., St. Louis, MO, Prod. No. E6888), or TGF $\beta$ 1 neutralizing antibody (R & D Systems) were added in the amounts indicated below to separate wells in triplicate to block the renin cascade at different sites after stimulation by renin:

10<sup>-5</sup> M Renin Inhibitor R042-5892 (Hoffman-LaRoche)

30 ng/ml Anti-TGF $\beta$ 1 antibody (R & D Systems, #AB 101 NA)

30 ng/ml Chicken IgG (control for anti-TGF $\beta$ 1 antibody, R & D Systems, # AB 101 C)

10<sup>-5</sup> M Enalapril<sup>™</sup> (Sigma Chemical Co., St. Louis, MO)

10<sup>-5</sup> M Losartan<sup>™</sup> (Merck Pharmaceuticals, West Point, PA)

These inhibitors were added at zero time with 10<sup>-7</sup> M human recombinant renin (Hoffman-LaRoche).

As shown in Figure 3, use of inhibitors that block renin's action to increase Angiotensin II, i.e. blocking Angiotensin I production from Angiotensinogen (Ro 42-5892), blocking Angiotensin I conversion to Angiotensin II (Enalapril<sup>™</sup>) and blocking binding of Angiotensin II to its type I receptor (Losartan<sup>™</sup>), does not reduce the renin-induced increase in TGF $\beta$  production. These results demonstrate for the first time an alternative pathway in which TGF $\beta$  production is stimulated by renin.

### EXAMPLE III

#### DEMONSTRATION OF INHIBITION OF TGF $\beta$ BY BLOCKING RENIN IN VIVO IN THE PRESENCE OF AN ANTI-FIBROTIC DRUG

5 In this example, a known fibrotic disease drug, Enalapril<sup>™</sup> which inhibits the production of Angiotensin II, is combined with an inhibitor of renin, antisense renin oligonucleotide, to obtain an enhanced therapeutic effect on fibrotic disease in an animal model.

10 Rats are administered Enalapril<sup>™</sup> in their drinking water prior to anti-thymocyte serum injection, e.g. three (3) days prior to injection. Anti-thymocyte antibody, e.g. OX-7, is injected intravenously into the rats at day three to produce fibrotic disease. (Bagchus et al., Lab. Invest. 55:680-687 (1986)). Renin antisense oligonucleotides are administered one hour following administration of OX-7 by introducing the oligonucleotides into a suitable vehicle, such as HVJ liposomes, and injecting the formulations into the left renal artery of Sprague Dawley rats as described for renin genes by Arai et al., *Biochem. And Biophys. Res. Comm.* 206(2):525-532 (1995), incorporated by reference herein. A control consisting of nonsense encoding oligonucleotides (e.g. derived from the renin antisense gene sequence) is also injected into the left renal artery of additional rats. The renin antisense localizes in the juxtaglomerular apparatus of the glomerulus where renin is produced blocking renin production.

15  
20  
25 Animals are sacrificed on day 7 and kidney tissue samples are taken for analysis of levels of TGF $\beta$  in the glomeruli. Glomeruli are sieved individually from each rat and placed in culture in suitable medium for three days. At the end of culture, culture supernatant is harvested by centrifugation and TGF $\beta$ , fibronectin and PAI-1 production are determined as markers of fibrotic renal disease severity. Other glomeruli are pooled and used to isolate RNA. RNA is used by standard methods to quantitate expression of mRNAs of interest, including TGF $\beta$ , fibronectin and collagens.

30 Glomeruli are also examined histologically for phenotypical changes, e.g. changes resulting from deposition for ECM. Phenotypic changes are associated with pathological

alteration of glomeruli indicative of fibrotic disease. Such changes include expansion of extracellular matrix in the mesangial area of the kidney in animal models and the presence of activated mesangial cells which have acquired the characteristics of fibroblasts, e.g. expressing  $\alpha$ -smooth muscle actin and interstitial collagen, indicating progressive glomerular injury (Johnson et al., *J. Am. Soc. Nephrol.* 2:S190-S197 (1992)). Tissue for light microscopy is fixed in formaldehyde, then dehydrated in graded ethanol and embedded in paraffin. Sections are cut at 3  $\mu$ m thickness and are stained with with the periodic Schiff reagent. The paraformaldehyde-fixed renal section of the rats are also incubated with mouse anti-human renin monoclonal antibody (Kaiichi Radioisotope Labs, Ltd., Tokyo, Japan), mouse anti- $\alpha$ -smooth muscle actin monoclonal antibody (Immunotech S. A. (Marseille, France) and rabbit anti-collagen antibodies (Chemicon, Temecula, CA, prod. No. AB755). The sections are further processed using Vectastain ABC Kit (Vector Laboratories, Inc., Burlingame, CA).

Results of antibody binding indicate the extent of glomerular injury and the effects of inhibition of renin on such injury.

#### EXAMPLE IV

##### ADDITIONAL DEMONSTRATION THAT RENIN UPREGULATES TGF $\beta$ IN HUMAN MESANGIAL CELLS

Primary cultures of adult human mesangial cells were grown from human nephrectomy tissues using standard methods. Cells were passaged 4-7 times and then plated (3,000 cell/cm<sup>2</sup>) in 12 well plates in 2ml of medium (Mesangial Basal Medium (Clonetics Corp.) containing 5% FCS, 10  $\mu$ g/ml penicillin and 100  $\mu$ g/ml streptomycin) and allowed to grow to 70% confluency for 48 hours at 37°C, 5% CO<sub>2</sub>. Cultures were washed three times using sterile phosphate buffered saline at room temperature and then 2 ml/well of serum free MBM medium to induce quiescence. After 48 hours, the serum-free medium was removed and 2 ml/well of fresh serum-free medium was added for 24 hours. Human recombinant renin (HrRenin, Hoffman-La Roche Ltd., Basel, Switzerland) in concentrations from 10<sup>-6</sup> to 10<sup>-12</sup> M was added to each well for 24 hours. A blank (no HrRenin) was used as a control. Cells and supernatants were harvested by

centrifugation after 24 hrs of culture and frozen at -70°C until analysis.

The total production and release of TGF $\beta$  into the culture supernatant was measured using an ELISA kit (R & D Systems). Induction of the matrix protein fibronectin (Fn) in the supernatant was measured using anti-fibronectin antibodies in an ELISA to provide further confirmation of induction of TGF $\beta$ . Renin-induced induction of TGF $\beta$ , fibronectin and PAI-1 mRNA were measured over time using semi-quantitative RT-PCR in a multiplex system where multiple cDNAs are amplified simultaneously according to Dostal et al., *Anal. Biochem.* 223:239-250 (1994), incorporated by reference herein. Determinations were done in triplicate mesangial cell cultures.

#### (1) Determination of Dose Dependency of Renin Induction of TGF $\beta$

As shown in Figure 4, statistically significant ( $p < 0.05$ ) dose dependent increases in TGF $\beta$  (Figure 4A) and Fn production (Figure 4B) were observed, peaking with 2- and 1.4-fold increases at  $10^{-6}$ M HrRenin, respectively. Time course experiments using  $10^{-7}$ M HrRenin revealed significant increases in TGF $\beta$  and Fn production at 24 and 48 hours ( $p < 0.03$  (Figure 5A and B). As shown in Figure 6A-C, renin-induced increases in TGF $\beta$ , PAI-1 and Fn mRNAs peaked at 4 hours with increases from 1.5- to 2-fold.

#### (2) Demonstration that Renin Upregulation of TGF $\beta$ is not mediated through Renin Enzymatic Activity or Angiotensin II

Renin inhibitor Ro42-5892 (Hoffman-LaRoche, Basel, Switzerland), Losartan<sup>™</sup> (Merck Pharmaceuticals, West Point, PA), Enalapril<sup>™</sup> (Sigma Chemical Co., St. Louis, MO, Prod. No. E6888), or TGF $\beta$ 1 neutralizing antibody (R & D Systems) were added in the amounts indicated below to separate wells in triplicate to block the renin cascade at different sites after stimulation by renin:

$10^{-5}$  M Renin Inhibitor R042-5892 (Hoffman-LaRoche)

$10^{-5}$  M Enalapril<sup>™</sup> (Sigma Chemical Co., St. Louis, MO)

$10^{-5}$  M Losartan<sup>™</sup> (Merck Pharmaceuticals, West Point, PA)

Controls included neutralizing antibody to TGF $\beta$  (ATG) and control IgG (TgG)

These inhibitors were added at zero time with  $10^{-7}$  M human recombinant renin (Hoffman-LaRoche).

As shown in Figure 7, use of inhibitors that block renin's action to increase Angiotensin II, i.e. blocking Angiotensin I production from Angiotensinogen (RO 42-5892), blocking Angiotensin I conversion to Angiotensin II (Enalapril<sup>™</sup>) and blocking binding of Angiotensin II to its type I receptor (Losartan<sup>™</sup>), does not reduce the renin-induced increase in TGF $\beta$  production.

These results provide additional evidence that renin upregulates TGF $\beta$  production by human mesangial cells through a mechanism which is independent of renin's enzymatic action to convert angiotensin to Angiotensin I, and independent of Angiotensin II generation. These results may have profound implications for progression of fibrotic renal disease, particularly in states of high plasma renin as are observed with therapeutic Angiotensin II blockade. Thus, the use of therapeutic agents such as Enalapril<sup>™</sup> or Losartan<sup>™</sup> for Angiotensin blockade may not be optimal as treatment agents because of resulting high renin levels, preventing a therapeutic reduction in TGF $\beta$ . In addition, antagonists developed to block the site on renin that acts in the Angiotensin II pathway, would not be expected to block the action of renin that is independent of this pathway. Therefore, effective therapy of fibrotic diseases must take these multiple pathways for TGF $\beta$  increase into consideration.

## EXAMPLE V

### DEMONSTRATION OF THE ABILITY OF tPA TO INCREASE PLASMIN DEGRADATION OF ACCUMULATED ECM IN VIVO

In this Example, recombinant tissue type plasminogen activator (rtPA) was shown to promote generation of the protease plasmin in nephritic glomeruli and to degrade pathological ECM proteins leading to a therapeutic reduction in matrix accumulation.



Six Sprague-Dawley rats with were injected with phosphate buffered saline (PBS, as a control) and 18 rats were injected with 300 ug of mouse monoclonal OX7 antibody produced in the laboratory using commercially obtained hybridoma cells (American Type Culture Collecton (Rockville, MD, USA; Peters et al., *Kidney Internatl.* 54:1570-1580 (1998)) on day 1 to induce anti-Thy-1 nephritis. Injection of the anti rat-thymocyte antibody intravenously causes binding to an epitope in rat glomerular mesangial cells call Thy 1.1. The complement-mediated mesangial cell lysis that follows initiates a cascade of tissue injury, followed by a repair process that involves induction of TGF $\beta$ -driven production and deposition of ECM components. In addition, the plasmin protease system is altered such that PA is decreased and PAI-1 is markedly increased. These alterations favor decreased plasmin generation which decreases matrix turnover and enhances matrix accumulation. Plasmin is the key to mesangial cell matrix turnover (Baricos et al, *Kidney Int.* 47:1037-1047 (1995)).

Three days after the initial injection, rtPA (Genentech, Inc., San Francisco, CA) in a formulation designed for rodent intravenous injection (GenBank EO8757) or PBS was injected intravenously. Injections were repeated twice a day from day 3 to day 5. RtPA was injected i.v. at a dose of 1 mg/kg BW (n=6). Controls received saline (n=6). Glomerular staining for ECM matrix proteins (collagen type I and III, fibronectin EDA+ and tenascin) and glomerular mRNA levels of TGF $\beta$ 1, fibronectin and PAI-1 were evaluated at day 6. Localization of rtPA in nephritic glomeruli and the effect of rtPA on glomerular plasmin were investigated. Rats were sacrificed at day 6 and kidney tissues excised, fixed in formalin and frozen for histological analysis.

Table 1

<u>Groups of Six Rats</u>	<u>Treatment</u>
Group 1-Normal controls	300 ug of PBS on day 1, then 300 ug PBS 2X
Group 2- Disease control	300 ug of OX7 on day 1, then 300 ug PBS 2X
Group 3 - Disease + Dose 1	300 ug of OX7 on day 1, then 0.25 mg/day rtPA 2X/day on days 3, 4 and 5

Kidney tissue sections were stained for extracellular matrix using Periodic Acid Schiff (PAS) using standard procedures and were stained for specific relevant matrix proteins such as Collagen I, Collagen IV, Fibronectin EDA and tenascin using standard immunohistochemical staining procedures. Matrix proteins were scored by image analysis of 30 glomeruli per rat.

Figure 8A (control) and B (tPA) show an overall decrease in matrix accumulated as a result of tPA treatment. Compared to the untreated, disease control group (Figure 9A-D), the percentage of the glomerular area with positive staining was significantly lower in the rtPA treated group at day 6 for fibronectin EDA+(FN) ( $19 \pm 2$  vs.  $14 \pm 1$ ,  $p < 0.01$ ), laminin ( $35 \pm 2$  vs.  $25 \pm 2$ ,  $p < 0.001$ ), type I collagen  $33 \pm 1$  vs.  $21 \pm 3$ ,  $p < 0.001$ ) and type IV collagen ( $27 \pm 2$  vs.  $23 \pm 1$ ,  $p < 0.01$ ). Glomerular levels of TGF $\beta$ 1, FN and PAI-1 mRNA were unchanged (Figure 10). rtPA co-localized with fibrin along the glomerular capillary loops and in the mesangium.

rtPA was injected into nephritic rats 10, 20 and 30 minutes before sacrifice. At sacrifice, glomeruli were isolated and placed in culture with a chromogenic substrate for tPA. Plasmin generation by nephritic glomeruli, as shown in Figure 11, was significantly elevated in tPA treated nephritic glomeruli compared to nephritic glomeruli from disease control rats.

This example demonstrates that injected rtPA binds fibrin in nephritic glomeruli where it increases plasmin generation and promotes pathological ECM degradation. rtPA may thus be used in the methods of the invention as an ECM degrading agent.

## EXAMPLE VI

### EFFECT OF ADMINISTRATION OF TGF $\beta$ INHIBITORY AGENTS AND AGENTS THAT PROMOTE DEGRADATION OF ECM

In this example, at least one agent that inhibits TGF $\beta$ , anti-TGF $\beta$  antibody or decorin, is administered in combination with an ECM degrading agent, such as rtPA to reduce excess ECM

accumulation and degrade accumulated ECM in an animal model of glomerulonephritis.

Sprague-Dawley rats are treated as described in the above Examples to induce nephritis. Groups of six (6) rats each include untreated disease controls, rats treated with tPA alone as in Example V, above, rats treated with Enalapril™ alone (200 mg/day) in drinking water and rats treated with both intravenous rtPA and Enalapril™ in drinking water. On day 6 rats are sacrificed and kidney sections are excised, fixed in formalin and frozen for histological analysis. Glomeruli are isolated and used for in vitro analysis of production of TGFβ, fibronectin and PAI-1 using ELISA assays of culture supernatants and for isolation of RNA for Northern analysis of message levels of TGFβ, fibronectin and PAI-1. Tissue samples are stained for ECM proteins and glomerular mRNA levels of TGFβ1, fibronectin and PAI-1.

It is expected that the results of treatments with both anti-TGFβ antibody and rtPA treatment are significantly lower positive staining both in PAS stained tissue and in glomeruli stained for specific matrix components, as shown in Example V, compared with groups treated with either agent alone or in the control disease group.

## EXAMPLE VII

### DEMONSTRATION OF THE EFFECTS OF ADMINISTRATION OF A PAI-1 MUTANT ON EXTRACELLULAR MATRIX DEGRADATION

The human PAI-1 mutant used in this experiment (see WO 97/39028) was constructed on the wild-type PAI-1 background (Ginsburg et al., *J. Clin. Invest.* 78:1673-1680 (1986)), and disabled by the introduction of two Arg residues at positions 333 and 335 of the mature protein, which are also referred to as residues P14 and P12 of the reactive center loop (Lawrence, *Adv. Exp. Med. Biol.* 425:99-108 (1997)). Upon interaction with a proteinase, these substitutions greatly retard the insertion of the reactive center loop into β-sheet A and prevent the mutant from adopting the latent conformation. Since loop insertion results in loss of vitronectin affinity (Lawrence et al., 1997, supra), the PAI-1 mutant retains significant vitronectin activity while

failing to inhibit all plasminogen activators.

Four to six week old male Sprague-Dawley rats (Sasco, Inc., Omaha, NE) were treated as described in the above Examples to induce anti-thy-1 nephritis by intravenous injection of the monoclonal anti-thymocyte antibody OX-7 350 mg/200 g body weight. Groups of six (6) rats included a normal control group (injected with saline), an untreated disease control group (injected with PBS), and a group treated with 1 mg/Kg PAI-1 mutant injected once a day beginning 24 hours after induction of ATS nephritis and ending at day 5. Two additional groups of rats were treated with 1) 100 mg/liter of Enalapril (in drinking water) with a loading dose of Enalapril given by gavage 24 hr after disease induction followed by 100 mg/liter of Enalapril in drinking water, and 2) a 6% low protein diet (Teklad, Madison, WI, diet number TD86551) started 24 hours following disease induction.

Rats were sacrificed at day 6 and kidney tissues excised, fixed in formalin and frozen for histological analysis. Kidneys were perfused in situ with cold buffered saline (PBS) at pH 7.4, and then excised. Pieces of cortex were removed and either snap frozen in 2-methylbutane that had been cooled in liquid nitrogen or fixed in 10% neutralized formalin for immunohistologic examination. The capsules were removed and the cortical tissue dissected out and minced with a razor blade prior to isolation of glomeruli by standard graded sieving. Kidney tissue sections were stained for extracellular matrix using Periodic Acid Schiff (PAS) using standard procedures and were stained for specific relevant matrix proteins such as Collagen I, Collagen IV, Fibronectin EDA and tenascin using standard immunohistochemical staining procedures. Matrix proteins were scored by a blinded observer. 20 glomeruli per rat were evaluated. Isolated glomeruli were also used to determine glomerular mRNA levels of TGF $\beta$ 1, fibronectin and PAI-1 at day 6.

Reagents to measure plasmin activity, including plasminogen, low molecular weight u-PA and H-D-Val-Leu-Lys-p-nitroanilide (S-2251) were obtained from KabiVitrum (Franklin, OH). PAI-1 activity was assayed by measuring the hydrolysis of synthetic substrate by formed plasmin in the presence of plasminogen (Marshall et al., *J. Biol. Chem.* 265:9198-8204 (1990)). Assays were performed in polyvinyl chloride microtiter plates. The total volume of 125  $\mu$ l was

comprised of the following: sample, H-D-Val-Leu-Lys-P-nitroanilide (0.01  $\mu$ M) and plasminogen (0.03  $\mu$ M) in 0.5% Triton X-100, 0.1 M Tris, at pH 8.0. The amount of p-nitroaniline released was measured at 410 nm with a Thermomax microplate reader (Molecular Devices, Menlo Park, CA). A standard curve was generated with each assay using low molecular weight human u-PA. Each sample was also assayed without plasminogen to establish the plasminogen-dependence of the enzyme activity. The plasmin activity in culture supernatant or cell lysate was expressed as IU/1000 glomeruli.

Figure 12 shows an increase in plasmin generation of glomeruli in culture as a result of injection of the PAI-1 mutant. Compared to the untreated, disease control group, the glomerular plasmin activity was significantly higher in the PAI-1 treated group, being approximately halfway between the activity of disease controls and normal glomeruli. Notably, the significant increase in glomerular plasmin activity in nephritic glomeruli was observed with the PAI-1 mutant 24 hours following the final injection.

In addition, treatment with the PAI-1 mutant resulted in decreased accumulation of Collagen Type I, relative to diseases controls (Figure 13), while glomerular levels of TGF $\beta$ 1, FN, PAI-1 mRNA and Collagen I mRNA were not significantly altered. The decreased accumulation of Collagen Type I together with the fact that the Collagen I mRNA does not significantly decrease suggests enhanced extracellular matrix degradation rather than decreased production of Collagen I.

These results suggest that the increase in glomerular plasmin activity with a PAI-1 mutant can be titrated to avoid large increases in plasmin generation that may lead to hemorrhaging. Thus, the dose of the PAI-1 mutant may be altered, for example by doubling the dose, to increase glomerular plasmin activity to normal, but not excessive, levels to decrease deleterious accumulation of extracellular matrix. In addition, the time of treatment may be extended, for example to 10 days to obtain desired degradation.

Various publications are cited herein that are hereby incorporated by reference in their entirety.

As will be apparent to those skilled in the art in which the invention is addressed, the present invention may be embodied in forms other than those specifically disclosed without departing from the spirit or potential characteristics of the invention. Particular embodiments of the present invention described above are therefore to be considered in all respects as illustrative and not restrictive. The scope of the invention is as set forth in the appended claims and equivalents thereof rather than being limited to the examples contained in the foregoing description.

5

## WE CLAIM:

1. A method for treating a condition associated with the excess accumulation of extracellular matrix in a tissue and/or organ or at a dermal wound site comprising:

a) reducing the excess accumulation of extracellular matrix associated with TGF $\beta$  overproduction and/or activity in an organ or tissue, or at a wound site; and

b) degrading excess accumulated extracellular matrix in said tissue and/or organ or wound site;

whereby the accumulation of extracellular matrix in said tissue and/or organ or wound site is reduced from the level existing at the time of treatment.

2. The method of claim 1 wherein the accumulation of extracellular matrix is reduced to a level which does not interfere with normal functioning of the tissue or organ or result in scarring.

3. The method of claim 1 wherein said step of reducing the accumulation of extracellular matrix comprises administering at least one agent that inhibits TGF $\beta$  in an amount sufficient to inhibit TGF $\beta$  overproduction and/or activity.

4. The method of claim 3, wherein said agent that inhibits TGF $\beta$  is selected from the group consisting of selected from the group consisting of inhibitors of aldosterone, inhibitors of angiotensin II, anti-TGF $\beta$  antibodies, renin, ACE inhibitors, AII receptor antagonists, proteoglycans and ligands for the TGF $\beta$  receptor.

5. The method of claim 4 wherein said agent is a proteoglycan selected from the group consisting of decorin, biglycan, fibromodulin, lumican, betaglycan and endoglin.

6. The method of claim 4 wherein said ACE inhibitor is Enalapril<sup>™</sup>.

7. The method of claim 4 wherein said AII receptor antagonist is Losartan<sup>tm</sup>.
8. The method of claim 1 wherein said step of reducing the accumulation of extracellular matrix associated with TGF $\beta$  activity comprises contacting renin with an anti-renin agent.
9. The method of claim 1 wherein said step of degrading excess accumulated extracellular matrix comprises contacting said matrix with at least one protease in an amount sufficient to degrade excess accumulated extracellular matrix to a level that does not impair the normal function of said tissue and/or organ or result in scarring.
10. The method of claim 9 wherein said protease is selected from the group consisting of serine proteases, metalloproteinases and protease combinations.
11. The method of claim 1 wherein said step of degrading accumulated extracellular matrix comprises administering an agent which increases the amount of active protease sufficient to degrade excess accumulated matrix to a level that does not impair the normal function of said tissue and/or organ or result in scarring.
12. The method of claim 11 wherein said protease is selected from the group consisting of serine proteases, metalloproteinases and protease combinations.
13. The method of claim 12 wherein said protease is plasmin and said agent which increases the amount of active plasmin is tPA.
14. The method of claim 12 wherein said protease is plasmin and said agent which increases the amount of active plasmin is a PAI-1 mutant.
15. The method of claim 1 wherein said condition associated with the excess accumulation of extracellular matrix is a fibrotic condition.



16. The method of claim 15 wherein said fibrotic condition is selected from the group consisting of glomerulonephritis, adult or acute respiratory distress syndrome (ARDS), diabetes, diabetic kidney disease, liver fibrosis, kidney fibrosis, lung fibrosis, post infarction cardiac fibrosis, fibrocystic diseases, fibrotic cancer, post myocardial infarction, left ventricular hypertrophy, pulmonary fibrosis, liver cirrhosis, veno-occlusive disease, post-spinal cord injury, post-retinal and glaucoma surgery, post-angioplasty restenosis, renal interstitial fibrosis, arteriovenous graft failure and scarring.

17. The method of claim 1 wherein said tissue or organ is selected from the group consisting of kidney, lung, liver, heart, arteries, skin and the central nervous system.

18. The method of claim 1 wherein said condition associated with the excess accumulation of extracellular matrix is scarring.

19. The method of claim 3 wherein said agent that inhibits TGF $\beta$  is nucleic acid encoding the agent.

20. The method of claim 9 wherein said protease is nucleic acid encoding a protease.

21. A method for treating a condition associated with the excess accumulation of extracellular matrix in a tissue and/or organ or at a wound site comprising:

a) administering at least one agent that inhibits TGF $\beta$  to reduce additional accumulation of extracellular matrix associated with TGF $\beta$  activity in a tissue and/or organ or at a wound site; and

b) administering at least one agent to degrade excess accumulated extracellular matrix; whereby the accumulation of extracellular matrix in said tissue and/or organ or at said wound site is reduced to a level which does not interfere with normal functioning of the tissue or organ in which said extracellular matrix accumulated.

22. The method of claim 21 wherein said agent that inhibits TGF $\beta$  and said agent to degrade excess accumulated extracellular matrix are administered concurrently.

23. The method of claim 21 wherein said agent that inhibits TGF $\beta$  and said agent to degrade excess accumulated extracellular matrix are administered sequentially.

24. The method of claim 21 wherein the agent that inhibits TGF $\beta$  is an anti-TGF $\beta$  specific agent that binds to and inhibits the activity of TGF $\beta$ .

25. The method of claim 21 wherein the agent that inhibits TGF $\beta$  is an agent selected from the group consisting of inhibitors of aldosterone, inhibitors of angiotensin II, anti-TGF $\beta$  antibodies, renin, ACE inhibitors, AII receptor antagonists, proteoglycans and ligands for the TGF $\beta$  receptor.

26. A method for treating or preventing a condition associated with the excess accumulation of extracellular matrix in a tissue and/or organ, or at a wound site, comprising administering a combination of agents in an amount sufficient to inhibit TGF $\beta$  activity and/or production to prevent or reduce the excess accumulation of extracellular matrix associated with TGF $\beta$  overproduction in a tissue and/or organ, or at a wound site.

27. The method of claim 26 wherein the agents to inhibit TGF $\beta$  comprise an agent that binds to and inhibits the activity of TGF $\beta$  and an agent selected from the group consisting of inhibitors of aldosterone, inhibitors of angiotensin II, inhibitors of renin, ACE inhibitors and AII receptor antagonists.

28. The method of claim 27 wherein said agent that binds to and inhibits the activity of TGF $\beta$  is selected from the group consisting of anti-TGF $\beta$  antibodies, proteoglycans and ligands for the TGF $\beta$  receptor.

29. The method of claim 28 wherein said agent that binds to and inhibits the activity of TGF $\beta$  is a proteoglycan selected from the group consisting of decorin, biglycan, fibromodulin, lumican, betaglycan and endoglin.

30. The method of claim 26 wherein said agents are nucleic acids encoding said respective agents.

5 31. The method of claim 26 wherein said agents are administered to reduce TGF $\beta$  overproduction prior to excess accumulation of extracellular matrix.

32. A method for preventing or reducing excess extracellular matrix accumulation in a tissue or organ or at a wound site comprising inhibiting the overproduction of TGF $\beta$  present in an organ  
10 or tissue or at a wound site to prevent

33. The method of claim 32 further comprising the step of degrading excess accumulated extracellular matrix in said tissue and/or organ or at said wound site.

15 34. The method of claim 32 wherein said accumulation of extracellular matrix is reduced to a level which does not interfere with the normal functioning of the tissue and/or organ in which said extracellular matrix accumulated or scarring is prevented or reduced.

20 35. A composition for reducing TGF $\beta$  overproduction in a tissue and/or organ or at a wound site comprising a combination of agents that inhibit TGF $\beta$  activity and/or production in a pharmaceutically acceptable carrier.

25 36. The composition of claim 35 wherein said agents comprise an agent that binds to and inhibits TGF $\beta$  and an agent that inhibits a pathway resulting in TGF $\beta$  production.

30 37. A composition for preventing or reducing excess accumulation of extracellular matrix associated with TGF $\beta$  activity comprising at least one agent that inhibits TGF $\beta$  activity and/or production and at least one agent that degrades excess accumulated extracellular matrix in a pharmaceutically acceptable carrier.

38. The composition of claim 37 wherein said agent that inhibits TGF $\beta$  is an agent that binds to and inhibits TGF $\beta$ , and said agent that degrades excess accumulated extracellular matrix is a protease.

39. The composition of claim 38 wherein said agent that inhibits TGF $\beta$  is selected from the group consisting of inhibitors of aldosterone, inhibitors of Angiotensin II, anti-TGF $\beta$  antibodies, renin, ACE inhibitors, AII receptor antagonists, proteoglycans and ligands for the TGF $\beta$  receptor.

40. The composition of claim 38 wherein said protease is selected from the group consisting of serine proteases, metalloproteinases and protease combinations.

41. The composition of claim 39 wherein said proteoglycan is selected from the group consisting of decorin, biglycan, fibromodulin, lumican, betaglycan and endoglin.

42. The composition of claim 37 wherein said agent that inhibits TGF $\beta$  and said agent that degrades excess accumulated extracellular matrix are nucleic acids encoding said agents respectively.

43. The composition of claim 37 wherein said agent that degrades excess accumulated extracellular matrix is an agent that increases the amount of active protease present in the tissue and/or organ or at a wound site.

44. The composition of claim 43 wherein said protease is plasmin and said agent is tPA.

45. The composition of claim 43 wherein said protease is plasmin and said agent is a PAI-1 mutant.

46. A composition for reducing the excess accumulation of extracellular matrix in a tissue and/or organ, or at a dermal wound site, comprising an agent that increases the amount of active protease present in the tissue and/or organ or at the wound site in a pharmaceutically acceptable carrier.

47. The composition of claim 46 wherein said protease is plasmin and said agent is tPA.
48. The composition of claim 46 wherein said protease is plasmin and said agent is a PAI-1 mutant.
- 5
49. A composition for treating a condition associated with the excess accumulation of extracellular matrix in a tissue and/or organ or at a dermal wound site comprising a combination of agents that inhibit TGF $\beta$  in a pharmaceutically acceptable carrier.
- 10
50. The composition of claim 49 wherein said agents that inhibit TGF $\beta$  are selected from the group consisting of inhibitors of aldosterone, inhibitors of inhibitors of aldosterone, inhibitors of angiotensin II, anti-TGF $\beta$  antibodies, renin, ACE inhibitors, AII receptor antagonists, proteoglycans and ligands for the TGF $\beta$  receptor.
- 15
51. The composition of claim 50 wherein said proteoglycans are selected from the group consisting of decorin, biglycan, fibromodulin, lumican, betaglycan and endoglin.




FIGURE 1

# Repin Increases TGF- $\beta$ Production by Cultured Human Mesangial Cells in a Dose-dependent Manner




FIGURE 2

# Effect of Blocking Agents on TGF- $\beta$ Production by Human-Mesangial Cells in Culture in Response to Stimulation by $10^{-7}M$ Renin




FIGURE 3







FIGURE 4





FIGURE 5

6/13




FIGURE 6




FIGURE 7

## Effect of tPA on anti-Thy-1 nephritis at day 6




FIGURE 8

09/869820, 070504

**Staining Index (%)**  
(percentage of positive staining area/glomerular area)




FIGURE 9

## Effect of tPA on glomerular mRNA expressions at Day 6




FIGURE 10

## Plasmin activity of isolated glomeruli





FIGURE 11



**PAI-1 Mutant treatment enhances  
glomerular plasmin generation**



**FIGURE 12**




FIGURE 13

## United States Patent Application

## COMBINED DECLARATION AND POWER OF ATTORNEY

As a below named inventor I hereby declare that: my residence, post office address and citizenship are as stated below next to my name; that

I verily believe I am the original, first and sole inventor (if only one name is listed below) or a joint inventor (if plural inventors are named below) of the subject matter which is claimed and for which a patent is sought on the invention entitled: METHODS FOR TREATING CONDITIONS ASSOCIATED WITH THE ACCUMULATION OF EXCESS EXTRACELLULAR MATRIX

The specification of which

- a. ☒ is attached hereto  
 b. ☐ was filed on \_\_\_\_\_ as application serial no. \_\_\_\_\_ and was amended on \_\_\_\_\_ (if applicable) (in the case of a PCT-filed application) described and claimed in international no. \_\_\_\_\_ filed \_\_\_\_\_ and as amended on \_\_\_\_\_ (if any), which I have reviewed and for which I solicit a United States patent.

I hereby state that I have reviewed and understand the contents of the above-identified specification, including the claims, as amended by any amendment referred to above.

I acknowledge the duty to disclose information which is material to the patentability of this application in accordance with Title 37, Code of Federal Regulations, § 1.56 (attached hereto).

I hereby claim foreign priority benefits under Title 35, United States Code, § 119/365 of any foreign application(s) for patent or inventor's certificate listed below and have also identified below any foreign application for patent or inventor's certificate having a filing date before that of the application on the basis of which priority is claimed:

- a. ☐ no such applications have been filed.  
 b. ☒ such applications have been filed as follows:

FOREIGN APPLICATION(S), IF ANY, CLAIMING PRIORITY UNDER 35 USC § 119			
COUNTRY	APPLICATION NUMBER	DATE OF FILING (day, month, year)	DATE OF ISSUE (day, month, year)
PCT	PCT/US00/00179	5 January 2000	
ALL FOREIGN APPLICATION(S), IF ANY, FILED BEFORE THE PRIORITY APPLICATION(S)			
COUNTRY	APPLICATION NUMBER	DATE OF FILING (day, month, year)	DATE OF ISSUE (day, month, year)

I hereby claim the benefit under Title 35, United States Code, § 120/365 of any United States and PCT international application(s) listed below and, insofar as the subject matter of each of the claims of this application is not disclosed in the prior United States application in the manner provided by the first paragraph of Title 35, United States Code, § 112, I acknowledge the duty to disclose material information as defined in Title 37, Code of Federal Regulations, § 1.56(a) which occurred between the filing date of the prior application and the national or PCT international filing date of this application.

U.S. APPLICATION NUMBER	DATE OF FILING (day, month, year)	STATUS (patented, pending, abandoned)

I hereby claim the benefit under Title 35, United States Code § 119(e) of any United States provisional application(s) listed below:

U.S. PROVISIONAL APPLICATION NUMBER	DATE OF FILING (Day, Month, Year)
60/114,795	5 January 1999

I hereby appoint the following attorneys and agents associated with Customer No. 26941, respectively and individually, as my attorneys and agents, with full power of substitution, to prosecute this application and to transact all business in the Patent and Trademark Office connected therewith.

Please direct all communication to the address associated with Customer No. 26941, which is currently:

SaraLynn Mandel  
Mandel & Adriano  
35 No. Arroyo Parkway, Suite 60  
Pasadena, California 91103

I hereby declare that all statements made herein of my own knowledge are true and that all statements made on information and belief are believed to be true; and further that these statements were made with the knowledge that willful false statements and the like so made are punishable by fine or imprisonment, or both, under Section 1001 of Title 18 of the United States Code and that such willful false statements may jeopardize the validity of the application or any patent issued thereon.

2	Full Name Of Inventor	Family Name Noble	First Given Name Nancy	Second Given Name A.
0	Residence & Citizenship	City Salt Lake City	State or Foreign Country Utah	Country of Citizenship US
1	Post Office Address	Post Office Address 4464 South Abinadi	City Salt Lake City	State & Zip Code/Country Utah, 84124/US
Signature of Inventor 201:			Date:	
2	Full Name Of Inventor	Family Name Border	First Given Name Wayne	Second Given Name A.
0	Residence & Citizenship	City Salt Lake City	State or Foreign Country Utah	Country of Citizenship US
2	Post Office Address	Post Office Address 4464 South Abinadi	City Salt Lake City	State & Zip Code/Country Utah, 84124/US
Signature of Inventor 202:			Date:	
3	Full Name Of Inventor	Family Name Lawrence	First Given Name Daniel	Second Given Name A.
0	Residence & Citizenship	City Derwood	State or Foreign Country Maryland	Country of Citizenship US <i>USX</i>
3	Post Office Address	Post Office Address 16525 Keala Terrace	City Derwood	State & Zip Code/Country Maryland, 20855/US
Signature of Inventor 203:			Date: 6/21/01	

§ 1.56 Duty to disclose information material to patentability.

(a) A patent by its very nature is affected with a public interest. The public interest is best served, and the most effective patent examination occurs when, at the time an application is being examined, the Office is aware of and evaluates the teachings of all information material to patentability. Each individual associated with the filing and prosecution of a patent application has a duty of candor and good faith in dealing with the Office, which includes a duty to disclose to the Office all information known to that individual to be material to patentability as defined in this section. The duty to disclose information exists with respect to each pending claim until the claim is canceled or withdrawn from consideration, or the application becomes abandoned. Information material to the patentability of a claim that is canceled or withdrawn from consideration need not be submitted if the information is not material to the patentability of any claims remaining under consideration in the application. There is no duty to submit information which is not material to the patentability of any existing claim. The duty to disclose all information known to be material to patentability is deemed to be satisfied if all information known to be material to patentability of any claim issued in a patent was cited by the Office or submitted to the Office in the manner prescribed by §§ 1.97(b)-(d) and 1.98. However, no patent will be granted on an application in connection with which fraud on the Office was practiced

I hereby appoint the following attorneys and agents associated with **Customer No. 26941**, respectively and individually, as my attorneys and agents, with full power of substitution, to prosecute this application and to transact all business in the Patent and Trademark Office connected therewith.

Please direct all communication to the address associated with **Customer No. 26941**, which is currently:

SaraLynn Mandel  
Mandel & Adriano  
35 No. Arroyo Parkway, Suite 60  
Pasadena, California 91103


I hereby declare that all statements made herein of my own knowledge are true and that all statements made on information and belief are believed to be true; and further that these statements were made with the knowledge that willful false statements and the like so made are punishable by fine or imprisonment, or both, under Section 1001 of Title 18 of the United States Code and that such willful false statements may jeopardize the validity of the application or any patent issued thereon.

2	Full Name Of Inventor	Family Name Noble	First Given Name Nancy	Second Given Name A
0	Residence & Citizenship	City Salt Lake City	State or Foreign Country Utah	Country of Citizenship US <i>USX</i>
1	Post Office Address	Post Office Address 4464 South Abinadi	City Salt Lake City	State & Zip Code/Country Utah, 84124/US
Signature of Inventor 201: <i>Nancy A Noble</i>			Date: <i>6/22/01</i>	
2	Full Name Of Inventor	Family Name Border	First Given Name Wayne	Second Given Name A
0	Residence & Citizenship	City Salt Lake City	State or Foreign Country Utah	Country of Citizenship US <i>USX</i>
2	Post Office Address	Post Office Address 4464 South Abinadi	City Salt Lake City	State & Zip Code/Country Utah, 84124/US
Signature of Inventor 202: <i>W A Border</i>			Date: <i>24 June 01</i>	
2	Full Name Of Inventor	Family Name Lawrence	First Given Name Daniel	Second Given Name A.
0	Residence & Citizenship	City Derwood	State or Foreign Country Maryland	Country of Citizenship US
3	Post Office Address	Post Office Address 16525 Keats Terrace	City Derwood	State & Zip Code/Country Maryland, 20855/US
Signature of Inventor 203:			Date:	

#### § 1.56 Duty to disclose information material to patentability.

(a) A patent by its very nature is affected with a public interest. The public interest is best served, and the most effective patent examination occurs when, at the time an application is being examined, the Office is aware of and evaluates the teachings of all information material to patentability. Each individual associated with the filing and prosecution of a patent application has a duty of candor and good faith in dealing with the Office, which includes a duty to disclose to the Office all information known to that individual to be material to patentability as defined in this section. The duty to disclose information exists with respect to each pending claim until the claim is canceled or withdrawn from consideration, or the application becomes abandoned. Information material to the patentability of a claim that is canceled or withdrawn from consideration need not be submitted if the information is not material to the patentability of any claim remaining under consideration in the application. There is no duty to submit information which is not material to the patentability of any existing claim. The duty to disclose all information known to be material to patentability is deemed to be satisfied if all information known to be material to patentability of any claim issued in a patent was cited by the Office or submitted to the Office in the manner prescribed by §§ 1.97(b)-(d) and 1.98. However, no patent will be granted on an application in connection with which fraud on the Office was practiced

United States Patent & Trademark Office  
Office of Initial Patent Examination -- Scanning Division



Application deficiencies found during scanning:

☐ Page(s) \_\_\_\_\_ of \_\_\_\_\_ were not present  
for scanning. (Document title)

☐ Page(s) \_\_\_\_\_ of \_\_\_\_\_ were not present  
for scanning. (Document title)

☐ Scanned copy is best available. *Drawing are very dark*